



U.S. Fish & Wildlife Service

# Mourning Dove

*Population Status, 2017*



## **Mourning Dove Population Status, 2017**

U.S. Fish and Wildlife Service  
Division of Migratory Bird Management  
Population and Habitat Assessment Branch  
11510 American Holly Drive  
Laurel, MD 20708-4002

August 2017

**Cover photograph:** David Dolton releasing a mourning dove at Chatfield State Park, Colorado. Photo by David Sharp.

David Dolton helped manage mourning doves until his retirement as the Western Webless Specialist for the Division of Migratory Bird Management, U.S. Fish and Wildlife Service.

### **Suggested citation:**

Seamans, M. E. 2017. Mourning dove population status, 2017. U.S. Department of the Interior, Fish and Wildlife Service, Division of Migratory Bird Management, Washington, D.C.

All Division of Migratory Bird Management reports are available on our web site at:  
<https://www.fws.gov/birds/surveys-and-data/reports-and-publications.php>

# MOURNING DOVE POPULATION STATUS, 2017

MARK E. SEAMANS, U.S. Fish and Wildlife Service, Division of Migratory Bird Management, 755 Parfet Street Suite 235, Lakewood, CO 80215

*Abstract:* This report summarizes information collected annually in the U.S. on survival, recruitment, abundance and harvest of mourning doves. Trends in the number of doves heard and seen per route from the all-bird Breeding Bird Survey (BBS) are reported, and absolute abundance estimates based on band recovery and harvest data are provided. Harvest and hunter participation are estimated from the Migratory Bird Harvest Information Program (HIP). BBS data suggested that the abundance of mourning doves over the last 51 years increased in the Eastern Management Unit (EMU) and decreased in the Central (CMU) and Western (WMU) Management Units. Estimates of absolute abundance are available only since 2003 and indicate that there were about 279 million doves in the U.S. as of 1 September 2016. Abundance (in millions of birds) varied among management units in 2016: EMU 62.3 (SE=3.5); CMU 171.0 (SE=13.6); and WMU 45.8 (SE=3.5). HIP estimates for mourning dove total harvest, active hunters, and total days afield in the U.S. in 2016 were 13,502,000 ± 557,500 (estimate ± SE) birds, 837,800 hunters, and 2,431,000 ± 96,900 days afield. Harvest and hunter participation at the management unit level were: EMU, 4,606,000 ± 216,700 birds, 303,100 hunters, and 789,600 ± 31,200 days afield; CMU, 7,334,600 ± 508,100 birds, 430,400 hunters, and 1,344,400 ± 90,500 days afield; and WMU, 1,561,400 ± 76,000 birds, 104,300 hunters, and 297,000 ± 14,900 days afield.

---

The mourning dove (*Zenaida macroura*) is one of the most abundant bird species in North America, and is familiar to millions of people. Authority and responsibility for management of this species in the U.S. is vested in the Secretary of the Interior. This responsibility is conferred by the Migratory Bird Treaty Act of 1918 which, as amended, implements migratory bird treaties between the U.S. and other countries. Mourning doves are included in the treaties with Great Britain (for Canada) and Mexico (U.S. Department of the Interior 2013). These treaties recognize sport hunting as a legitimate use of a renewable migratory bird resource.


Maintenance of dove populations in a healthy, productive state is a primary management goal. Management activities include population assessment, harvest regulation, and habitat management. Each year, tens of thousands of doves are banded and thousands of wings from harvested doves are analyzed to estimate annual survival, harvest rates, recruitment, and abundance. The resulting information is used by wildlife managers in setting annual hunting regulations (USFWS 2017). Past federal frameworks for hunting in the U.S. are in Appendix A.

## DISTRIBUTION

Mourning doves breed from southern Canada throughout the U.S. into Mexico, Bermuda, the Bahamas and Greater Antilles, and in scattered locations in Central America (Peterjohn et al. 1994, Fig. 1). Although mourning doves winter throughout much of their breeding range, the majority winter in the southern U.S., Mexico, and south through Central America to western Panama (Aldrich 1993, Mirarchi and Baskett 1994).

## POPULATION MONITORING

Within the U.S., three zones contain mourning dove populations that are largely independent of each other (Kiel 1959; Fig. 2). These zones encompass the principal breeding, migration, and U.S. wintering areas for each population. As suggested by Kiel (1959), these three zones were established as separate management units in 1960 (Kiel 1961). Since that time, management decisions have been made within the boundaries of the Eastern (EMU), Central (CMU),



**Figure 1.** Breeding and wintering ranges of the mourning dove (adapted from Mirarchi and Baskett 1994).

and Western (WMU) Management Units (Fig. 2). The EMU was further divided into two groups of states for some analyses. States permitting dove hunting were combined into one group (hunt) and those prohibiting dove hunting into another (non-hunt). Additionally, some states were grouped to increase sample sizes. Maryland and Delaware were combined; Vermont, New Hampshire, Maine, Massachusetts, Connecticut, and Rhode Island were combined to form a New England group. Even though Rhode Island is a hunt state, due to its small size and geographic location its data was included in this non-hunt group of states for analysis.

## Breeding Bird Survey

The North American Breeding Bird Survey (BBS; Robbins et al. 1986) is completed in June and is based on routes that are 24.5 miles long. Each route consists of 50 stops or point count locations at 0.5-mile intervals. At each stop, a 3-minute count is conducted whereby every bird seen within a 0.25-mile (400 m)

radius or heard is recorded. Surveys start one-half hour before local sunrise and take about 5 hours to complete. Data for birds heard and seen at stops are combined for BBS analyses.

Although the BBS is not used to inform annual mourning dove harvest management decisions, it is still of interest because it provides independent estimates of trends in abundance. Consequently, the 1966–2016 BBS trend information is included in this report. Current-year BBS data are not available in time for inclusion in the report.


## Banding Program

A national banding program was initiated in 2003 to improve our understanding of mourning dove population biology and to help estimate the effect of harvest on mourning dove populations. Doves are banded in July and August in most of the lower 48 states. Band recoveries occur almost exclusively during the U.S. hunting seasons which occur primarily between 1 September and 15 January (Appendix A).

Banding goals for each state (specified by Bird Conservation Region [BCR]) are based on a power analysis that estimated sample sizes necessary to achieve a desired precision in estimates of population growth rate at the management unit level (Otis 2009). A weighting factor based on the median BBS index during 1966–2008 was used to determine banding goals for each state within the management units. Within states, the amount of area in each BCR and associated median BBS indices were used to determine sample size allocation. Placement of banding stations is left to the judgment of the state banding coordinator.

## Harvest Survey

Wildlife professionals have long recognized that reliable harvest estimates are needed to monitor the impact of hunting. The U.S. Fish and Wildlife Service (USFWS), in collaboration with State agencies, initiated the Migratory Bird Harvest Information Program (HIP) in 1992. HIP became fully operational on a national scale in 1999. HIP is designed to enable the USFWS to conduct nationwide surveys that provide reliable annual estimates of the harvest of mourning doves and other migratory game bird species at state, management unit, and national levels.



**Figure 2.** Mourning dove management units with 2016-17 hunt and non-hunt states.

Under HIP, states provide the USFWS with the names and addresses of all licensed migratory bird hunters each year. Surveys are then sent to a sample of those hunters to estimate harvest and hunter participation (i.e., number of active hunters, total days afield) in each state. All states except Hawaii participate in the program.

### Parts Collection Survey

Age of individual doves can be determined by examination of their wings (Ruos and Tomlinson 1967, Braun 2014). Mourning dove wings are easily obtained during the hunting season and provide estimates of recruitment (number of young per adult in the population), which can be used to inform harvest management. From 2005–2009 some states collected wings for use in estimating age ratios in the fall populations. In 2007, the USFWS initiated the national Mourning Dove Parts Collection Survey, which expanded the geographical scope of the earlier state-based surveys.

The survey design for mourning dove wing collection follows that of waterfowl (Raftovich et al. 2016). The sampling frame is defined by hunters who identify themselves as dove hunters when purchasing a state hunting license and who were active dove hunters the previous year.

Each year, state and federal biologists classify wings during a 2-day wingbee hosted by the Missouri Department of Conservation in Lee’s Summit, Missouri. Wings of harvested mourning doves are classified as juveniles (hatch-year birds or HY) or adults (after-hatch-year birds or AHY). A significant portion of wings are classified as unknown age where molt has progressed to a late stage. These harvest age ratios (HY/AHY) are used to estimate recruitment (population age ratio) after accounting for uncertainty related to unknown-age wings and age-specific vulnerability to harvest (Miller and Otis 2010).

### Call-count Survey

The Mourning Dove Call-count Survey (CCS) was conducted from 1966 to 2013. The CCS was developed to provide an annual index of abundance specifically for mourning doves (Dolton 1993). The CCS was discontinued because the harvest strategy adopted for mourning doves in 2013 does not make use of data from the CCS, but rather relies on absolute abundance estimates. However, state and federal biologists are conducting a national study using a subset of the historical CCS routes to determine if point count surveys that use distance sampling methods (Buckland et al. 2001) can produce absolute abundance estimates that can be used to make regulatory decisions. Those interested in historic CCS information can access previous status reports for

mourning doves (available online at <https://www.fws.gov/birds/surveys-and-data/reports-and-publications.php>).

## METHODS

### Estimation of Trends in Abundance Indices

BBS trends were estimated using a log-linear hierarchical model and Bayesian analytical framework (Sauer et al. 2008, 2010). The hierarchical model has a rigorous and sound theoretical basis and the indices and trends are directly comparable because trends are calculated directly from the indices.

With the hierarchical model, the log of the expected value of the counts is modeled as a linear combination of stratum-specific intercepts and trends, a random effect for each unique combination of route and observer, a year effect, a start-up effect on the route for first year counts by new observers, and overdispersion (unexplained variation). Most of the parameters of interest are treated as random effects and some parameters are hierarchical in that they are assumed to follow distributions that are governed by additional parameters. The model is fit using Bayesian methods. Markov-chain Monte Carlo methods are used to iteratively produce sequences of parameter estimates which can be used to describe the distribution of the parameters of interest. Once the sequences converge, medians and credible intervals (CI, Bayesian confidence intervals) for the parameters are determined from the subsequent replicates. Annual indices are defined as exponentiated year and trend effects, and trends are defined as ratios of the year effects at the start and end of the interval of interest, taken to the appropriate power to estimate a yearly change (Sauer et al. 2008). Trend estimates are expressed as the average percent change per year over a given time period, while indices are expressed as the number of doves heard and seen per route.

Annual indices were calculated at the state, region (group of states), and dove management unit levels. Short- (recent 10-year period) and long-term (all years with data) trends were evaluated for each area. The median and 95th percentile credible intervals are presented for estimates. The extent to which trend credible intervals exclude zero can be interpreted as

the strength of evidence for an increasing or decreasing trend. Thus, there is evidence of a positive trend if the lower bound of the CI  $> 0$  and there is evidence of negative trend if the upper bound of the CI  $< 0$ . If the CI contains 0, then there is inconclusive evidence about trend in abundance. The reported sample sizes are the number of routes or sites on which trend estimates are based, which includes any route on which mourning doves were ever encountered in the region. BBS results are presented in Table 1.

### Estimation of Survival, Harvest, and Recruitment Rates, & Absolute Abundance

Band recovery models were used to estimate annual survival. A Seber parameterization (Seber 1970) with all dead recoveries was used to estimate survival rates. No adjustment was made to account for band reporting probabilities when estimating survival, and both direct and indirect recoveries were used. Only direct recoveries (bands recovered the same season applied) were used to estimate harvest rates and data were adjusted for band reporting rate (Sanders and Otis 2012) prior to analysis.

Age specific harvest and survival rates were estimated by state and management unit. Most states lacked sufficient sample sizes of banded birds to estimate annual survival rates; therefore, data were pooled over years to obtain mean annual estimates. Harvest rate for a year in a given state was only estimated when the number of banded birds in an age-class was  $>100$ . Harvest rates for Management Units were based on state-weighted harvest rate estimates. Each state's weight was the product of its habitat area (area within state presumed to be dove habitat) and average dove abundance estimated by the Call Count Survey index of doves heard 2009-2013 (the Call Count Survey was discontinued after 2013).

For estimating survival rates a model was formulated that allowed recovery rate to vary by state with an additive age effect, and allowed survival to vary by state and age. This model was used for inference regarding age and state specific survival rates.

The approach of Miller and Otis (2010) was used to estimate annual recruitment rates. Samples were limited to wings collected during the first two weeks of September to minimize the proportion of unknown

age wings and maximize the proportion of local birds in samples. Unknown age wings were assigned to an age-class based on previously estimated probabilities that adults will be in late stages of molt. Band recovery data was used to adjust age-ratio estimates for differential vulnerability to harvest.

A simple Lincoln-type estimator was used to estimate abundance from annual harvest and harvest rates (Otis 2006). Abundance for each year was estimated at the management unit level separately for juvenile and adult doves by dividing age-specific total harvest (from the USFWS Harvest Information Program [Table 3] and Parts Collection Survey [Table 6]) by age-specific harvest rates estimated from direct (first hunting season) recoveries of banded birds.

## RESULTS

### Breeding Bird Survey

*Eastern Management Unit.*—The BBS provided evidence that dove abundance increased in the EMU hunt and non-hunt states during the last 51 years (Table 1). Over the last 10 years there was evidence that abundance remained unchanged in the EMU hunt states and non-hunt states, and remained unchanged in the entire EMU.

*Central Management Unit.*—In the CMU, the BBS provided evidence that doves decreased in abundance over the last 51 years, but not the most recent 10 years (Table 1).

*Western Management Unit.*—The BBS provided evidence that dove abundance decreased in the WMU over the last 51 years but not during the most recent 10 years (Table 1).

### Harvest Survey

Preliminary results of mourning dove harvest and hunter participation from HIP for the 2015-16 and 2016-17 hunting seasons are presented in Tables 2 and 3, respectively. Current (2016-17) HIP estimates indicate that in the U.S. about 13.5 million mourning doves were harvested by about 840,000 hunters who spent about 2.4 million days afield. The EMU and CMU total harvest represented 34% and 54%, respectively, of the national harvest of doves while the

WMU represented 12% (Table 3). Mourning dove harvest and hunter participation were similar between the 2015-16 and 2016-17 seasons within each management unit and nationwide (Fig. 3, Tables 2 and 3).

Additional information about HIP, survey methodology, and results can be found in annual reports located at: <https://www.fws.gov/birds/surveys-and-data/reports-and-publications/hunting-activity-and-harvest.php>.

### Survival and Harvest Rates


Over the past 14 years 268,167 doves were banded in the EMU, 225,912 in the CMU, and 102,474 in the WMU during July and August (Table 4). There have been 17,057, 12,386, and 4,317 recoveries of banded birds in the EMU, CMU, and WMU, respectively.

Mean annual HY survival was similar between the management units (Table 5). AHY survival was similar in the CMU and WMU, but slightly lower in the EMU.

Mean annual harvest rate was higher for HY individuals compared to AHY individuals in all the management units (Fig. 3, Table 5). This relationship was more pronounced in the EMU (HY harvest rate 45% greater than AHY harvest rate) than the CMU (27% greater) and WMU (19% greater). Mean annual harvest rates by age-class (HY and AHY) were greater in the EMU than in the other management units (Table 5). Within the EMU, the harvest rate of birds banded in the North Atlantic states (predominantly non-hunt states) was much lower than that of the hunt states (Table 5).

### Recruitment


A total of 178,961 wings were obtained during 2007–2016 from birds harvested prior to September 15<sup>th</sup>. Overall recruitment rates were highest in the east and northwest and lowest in the Great Plains states and the southwest (Table 6). At the management unit level, the EMU had higher average annual recruitment and more annual variation compared to the CMU and WMU (Fig. 4). In 2016 the EMU and WMU experienced a lower-than-average age ratio in the fall



**Figure 3.** Estimated harvest (▲) and harvest rates of mourning dove 2003–2016. Harvest rates presented separately for hatch-year (□) and after-hatch-year (○).

population, whereas the CMU was near its long-term average (Table 6).

Mean population age ratios for all states and years are provided in Table 6. There was much variation in the sample sizes for individual states. However, sample sizes are sufficient to calculate precise estimates of recruitment for all states.




**Figure 4.** Estimated mourning dove fall population age ratios for each management unit, 2007–2016. Error bars represent 95% confidence intervals.

Age ratios for Florida are not estimated because hunting seasons there do not start until late September each year. At this late date most wings cannot be aged due to molt progression, precluding accurate estimates of age ratio.

### Absolute Abundance

Estimates of absolute abundance are available since 2003 (Fig. 5, Table 7). Estimates during the first 1 or 2 years may be biased in association with startup of the national mourning dove banding program when coordinators were gaining experience, and some states were not yet participants. In addition, age ratio information was not available for the first 4 years (the annual averages from later years were used for estimating abundance during this period). The most recent estimates indicate that there were 279 million mourning doves in the U.S. immediately prior to the 2016 hunting season. Compared to 2015, abundance appeared to increase in the WMU in 2016. Abundance appeared to remain relatively unchanged in the EMU and CMU.



**Figure 5.** Estimates and 95% confidence intervals of mourning dove absolute abundance by management unit and year, 2003–2016. Estimates based on band recovery and harvest data.

## ACKNOWLEDGMENTS

State wildlife agencies and the U.S. Fish and Wildlife Service (USFWS) cooperated to collect the data presented in this report. The following participated in the November 2016 Dove Wingbee: J. Odell (AZ), K. Fothergill (CA), R. Schultheis (KS), A. Friesen (KS),

V. Cikanek (KS), C. Powell (KY), J. Duguay (LA), J. Berdeen (MN), R. Bredesen (MO), L. Fendrick (OH), J. Neal (OK), M. Frisbie (TX), J. Timmons (TX), R. Rau (USFWS), S. Kelly (USFWS), T. Edwards (USFWS), and M. Seamans (USFWS). A special thanks to R. Bredesen (MO) for providing the space at the J.A. Reed Memorial Wildlife Area for the Wingbee. J. Sauer (USGS) analyzed the BBS data and provided statistical support. His commitment to the annual assessment of abundance data, report contributions, and extraordinary work hours during report preparation is highly valued. K. Fleming and B. Raftovich (USFWS) provided HIP and Parts Collection data, while T. Ceaser II, L. Heckstall, and P. Mathias (USFWS) entered data from the Dove Wingbee. M. Rogosky (USGS BBL) provided band and encounter data. P. Devers, J. Dubovsky, P. Garrettson, P. Padding, and R. Rau (USFWS) reviewed a draft of this report. V. Neatrou helped format and produce the cover page of this report. This report would not be possible without the significant contributions of all involved.

## LITERATURE CITED

Aldrich, J.W. 1993. Classification and distribution. Pages 47-54 in T.S. Baskett, M.W. Sayre, R.E. Tomlinson, and R.E. Mirarchi, Editors. Ecology and management of the mourning dove. Stackpole Books, Harrisburg, Pennsylvania, USA.

Braun, C.E. 2014. Use of secondary feathers to age mourning doves. *North American Bird Bander* 39:1–6.

Buckland, S.T., D.R. Anderson, K.P. Burnham, J.L. Laake, D.L. Borchers, and L. Thomas. 2001. Introduction to distance sampling. Oxford University Press Inc., New York.

Dolton, D.D. 1993. The call-count survey: historic development and current procedures. Pages 233–252 in T.S. Baskett, M.W. Sayre, R.E. Tomlinson, and R.E. Mirarchi, editors. Ecology and management of the mourning dove. Stackpole Books, Harrisburg, Pennsylvania, USA.

Kiel, W.H. 1959. Mourning dove management units, a progress report. U.S. Fish and Wildlife Service, Special Scientific Report—Wildlife 42.

Kiel, W.H. 1961. The mourning dove program for the future. *Transactions of the North American Wildlife and Natural Resources Conference* 26:418–435.

- Miller, D.A., and D.L. Otis. 2010. Calibrating recruitment estimates for mourning doves from harvest age ratios. *Journal of Wildlife Management* 74:1070–1079.
- Mirarchi, R.E. and T.S. Baskett. 1994. Mourning dove (*Zenaida macroura*). In A. Poole and F. Gill, editors, *The birds of North America*, No. 117. The Academy of Natural Sciences, Philadelphia and The American Ornithologists' Union, Washington, D.C., USA.
- Otis, D.L. 2006. A mourning dove hunting regulation strategy based on annual harvest statistics and banding data. *Journal of Wildlife Management* 70:1302–1307.
- Otis, D.L. 2009. Mourning dove banding needs assessment. U.S. Fish and Wildlife Service. Unpublished report. 22pp. Available online: <https://www.fws.gov/birds/surveys-and-data/webless-migratory-game-birds/doves-and-pigeons.php>
- Peterjohn, B. G., J. R. Sauer and W. A. Link. 1994. The 1992 and 1993 summary of the North American breeding bird survey. *Bird Populations* 2:46–61.
- Raftovich, R.V., S.C. Chandler, and K.A. Wilkins. 2016. Migratory bird hunting activity and harvest during the 2014-15 and 2015-16 hunting seasons. U.S. Fish and Wildlife Service, Laurel, Maryland, USA.
- Robbins, C.S., D. Bystrak, and P.H. Geissler. 1986. *The Breeding Bird Survey: its first fifteen years, 1965-1979*. U.S. Fish & Wildlife Service, Research Publication 157.
- Ruos, J. L., and R. E. Tomlinson. 1967. Results of mourning dove wing collection in the eastern management unit, 1966–67. U.S. Bureau of Sport Fisheries and Wildlife Administration Report, Washington, D.C., USA.
- Sanders, T. A., and D. L. Otis. 2012. Mourning dove reporting probabilities for web-address versus toll-free bands. *Journal of Wildlife Management* 76:480–488.
- Sauer, J. R., W. A. Link, W. L. Kendall, and D. D. Dolton. 2010. Comparative analysis of mourning dove population change in North America. *Journal of Wildlife Management* 74:1059–1069.
- Sauer, J. R., W. A. Link, W. L. Kendall, J. R. Kelly, and D. K. Niven. 2008. A hierarchical model for estimating change in American woodcock populations. *Journal of Wildlife Management*. 58:204–214.
- Seber, G.A.F. 1970. Estimating time-specific survival and reporting rates for adult birds from band returns. *Biometrika* 57:313–318.
- U.S. Department of the Interior. 2013. Final Supplemental Environmental Impact Statement: Issuance of annual regulations permitting the sport hunting of migratory birds. U.S. Fish and Wildlife Service. Washington, D.C., USA. Available online at: <https://www.fws.gov/birds/policies-and-regulations/regulations/how-regulations-are-set-the-process.php>
- U.S. Fish and Wildlife Service. 2017. Mourning Dove Harvest Strategy. U.S. Department of the Interior, Fish and Wildlife Service, Division of Migratory Bird Management, Washington, D.C. Available online at: <https://www.fws.gov/birds/surveys-and-data/webless-migratory-game-birds/doves-and-pigeons.php>

**Table 1.** Estimated trend<sup>a</sup> (percent change per year and lower and upper 95% credible intervals) in mourning dove abundance based on Breeding Bird Survey data for management units and states during 51-year (1966–2016) and 10-year (2007–2016) periods.

Management Unit State	51 year				10 year			
	N	Trend	Lower	Upper	N	Trend	Lower	Upper
Eastern	1,768	0.3	0.2	0.5	1,466	-0.2	-0.5	0.1
Hunt states	1,434	0.3	0.2	0.4	1,197	-0.3	-0.6	0.1
AL	101	-1.1	-1.5	-0.6	87	-1.0	-2.1	0.1
DE-MD	86	-0.1	-0.4	0.1	72	-0.5	-1.5	0.4
FL	99	2.1	1.4	2.8	79	0.8	-0.7	2.3
GA	104	-0.5	-0.9	-0.2	91	-0.4	-1.2	0.6
IL	102	0.6	-0.1	1.2	100	-3.3	-4.4	-2.2
IN	63	-0.3	-0.8	0.2	54	-2.2	-3.7	-0.8
KY	56	0.8	0.3	1.2	37	1.2	-0.1	2.7
LA	94	2.5	2	3.1	71	2.8	1.3	4
MS	54	0.1	-0.5	0.6	43	0.6	-0.7	2.3
NC	95	0.2	-0.2	0.5	81	0.3	-0.6	1.2
OH	78	0.5	-0.1	1.1	59	-0.1	-1.6	1.3
PA	127	0.9	0.5	1.4	101	-0.2	-1.2	0.9
SC	45	-0.1	-0.6	0.4	38	-0.2	-1.7	1
TN	29	-0.4	-0.9	0.1	23	0.0	-1.2	1.3
VA	59	-0.2	-0.6	0.2	51	0.0	-1.1	0.9
WI	95	1.3	0.8	1.8	90	-0.2	-1.5	1.1
WV	57	3.3	2.4	4.3	49	-0.3	-2.3	1.6
Non-hunt states	420	0.9	0.6	1.1	341	0.4	-0.3	1.1
MI	90	0.7	0.3	1.2	71	-0.6	-2	0.6
New England <sup>b</sup>	166	1.6	1.2	2.1	136	0.4	-0.8	1.6
NJ	42	-0.7	-1.3	-0.1	29	-0.6	-2	0.5
NY	126	1.5	1	2	104	1.0	-0.3	2.3
Central	1,205	-0.4	-0.5	-0.2	1,056	0.4	0	0.8
AR	55	0.7	-0.1	1.5	49	1.6	-0.5	3.7
CO	144	-0.6	-1.2	-0.1	132	-2.5	-4	-1.1
IA	38	0.8	0.3	1.4	32	1.1	-0.4	2.8
KS	65	-0.1	-0.6	0.4	62	0.0	-1.5	1.5
MN	78	-1.1	-1.6	-0.7	73	-0.6	-1.9	0.7
MO	89	-0.6	-1	-0.1	74	0.5	-0.6	1.9
MT	81	-0.7	-1.3	-0.1	71	0.7	-1.1	2.8
NE	51	-0.2	-0.7	0.3	46	0.2	-1	1.3
NM	80	-0.4	-1.2	0.4	60	0.2	-1.4	1.9
ND	50	-0.4	-1.2	0.4	47	-0.2	-1.8	1.3
OK	60	-1.2	-1.7	-0.7	53	-1.2	-2.5	0.2
SD	58	-0.1	-0.7	0.5	51	1.0	-0.6	2.8
TX	230	-0.4	-0.7	0	206	1.5	0.6	2.5
WY	126	-1.0	-1.8	-0.3	100	-0.6	-2.1	0.9
Western	716	-1.1	-1.6	-0.7	560	-0.7	-1.6	0.2
AZ	87	-1.2	-2.1	-0.4	64	-2.4	-4.2	-0.6
CA	251	-0.7	-1.3	-0.1	190	0.6	-0.9	2
ID	50	-0.7	-1.7	0.2	43	2.5	0.3	4.9
NV	45	-1.6	-3.5	0.2	31	-1.3	-4.7	2.1
OR	116	-1.0	-2.1	0.2	90	2.5	0.1	5
UT	102	-1.9	-2.7	-1.1	90	-2.1	-3.9	-0.3
WA	77	-0.1	-0.7	0.5	64	0.2	-1.2	1.7

<sup>a</sup>Trend estimated from annual indices derived from a log-linear hierarchical model fit using Bayesian methods. There is evidence of a positive trend if the lower CI > 0 and there is evidence of negative trend if the upper CI < 0. If the CI contains 0, then there is inconclusive evidence about trend in abundance.

<sup>b</sup> New England consists of CT, ME, MA, NH, RI, and VT; RI is a hunt state but was included in this group for purposes of analysis.

**Table 2.** Preliminary estimates and 95% confidence intervals (CI, expressed as the interval half width in percent) of mourning dove harvest and hunter activity for management units and states during the 2015 hunting season<sup>a</sup>.

Management Unit	Harvest		Active hunters		Hunter days afield		Harvest per hunter <sup>b</sup>	
	Estimate	CI	Estimate	CI	Estimate	CI	Estimate	CI
Eastern	4,644,900	8	297,000	† <sup>c</sup>	780,400	8	†	†
AL	428,000	19	26,700	12	59,500	17	16.1	23
DE	24,900	29	1,300	30	3,700	32	19.5	39
FL	141,900	39	7,000	34	22,400	29	20.3	52
GA	725,700	16	41,800	13	104,400	21	17.3	20
IL	283,600	30	18,400	18	55,800	38	15.4	36
IN	93,600	23	7,900	22	24,600	39	11.9	31
KY	286,500	29	15,200	21	38,300	26	18.9	35
LA	214,100	42	16,400	27	39,000	40	13.1	51
MD	63,100	28	5,200	23	12,300	24	12.1	36
MS	257,100	18	16,200	17	34,900	21	15.9	25
NC	734,300	29	48,700	20	117,500	25	15.1	35
OH	131,200	35	8,600	25	28,000	33	15.3	42
PA	119,200	35	17,800	28	58,500	41	6.7	44
RI	1,100	125	300	65	1,200	131	4.3	146
SC	548,600	24	27,900	19	72,000	21	19.7	31
TN	288,400	45	12,000	41	35,100	45	24.0	61
VA	229,500	20	17,300	16	40,400	27	13.2	25
WI	60,400	63	7,100	33	29,400	46	8.5	71
WV	13,700	21	1,500	13	3,700	26	9.3	30
Central	7,180,300	9	369,800	†	1,235,000	10	†	†
AR	252,400	22	17,800	24	37,600	22	14.2	33
CO	204,500	22	14,200	15	38,900	23	14.4	26
IA	111,500	18	9,200	15	24,600	16	12.1	23
KS	558,200	20	28,600	13	86,400	18	19.5	24
MN	96,700	86	9,700	48	28,200	54	10.0	100
MO	307,400	24	22,500	14	54,300	17	13.6	27
MT	18,000	54	1,600	49	5,100	54	11.0	69
NE	160,600	17	9,000	17	25,500	18	17.9	25
NM	111,900	22	7,000	11	23,100	14	16.0	25
ND	73,500	25	4,200	23	12,800	25	17.3	34
OK	294,000	18	18,200	15	45,300	17	16.1	24
SD	84,500	30	5,300	15	16,000	25	16.0	34
TX	4,892,100	13	220,700	11	834,000	14	22.2	18
WY	14,900	28	1,700	23	3,300	30	8.9	40
Western	1,332,200	8	82,000	†	226,500	9	†	†
AZ	401,400	7	17,100	3	53,900	5	23.5	8
CA	686,800	13	43,600	10	114,100	14	15.8	16
ID	100,700	45	5,200	26	18,500	40	19.3	52
NV	22,400	23	2,200	27	5,000	24	10.1	31
OR	22,500	35	3,200	31	9,100	43	7.0	48
UT	54,800	39	6,500	21	14,600	30	8.5	44
WA	43,600	41	4,200	28	11,200	33	10.3	49
United States	13,157,300	6	748,800	†	2,241,900	6	†	†

<sup>a</sup>Hunter number estimates at the management unit and national levels may be biased high, because the HIP sample frames are state specific; therefore hunters are counted more than once if they hunt in >1 state. Variance is inestimable.

<sup>b</sup>Seasonal harvest per hunter.

<sup>c</sup>No estimate available.

**Table 3.** Preliminary estimates and 95% confidence intervals (CI, expressed as the interval half width in percent) of mourning dove harvest and hunter activity for management units and states during the 2016 hunting season<sup>a</sup>.

Management Unit	Harvest		Active hunters		Hunter days afield		Harvest per hunter <sup>b</sup>	
	Estimate	CI	Estimate	CI	Estimate	CI	Estimate	CI
Eastern	4,606,000	9	303,100	† <sup>c</sup>	789,600	8	†	†
AL	396,000	21	31,400	14	65,400	17	12.6	25
DE	20,100	35	1,600	27	3,200	27	12.3	44
FL	88,200	43	7,600	37	18,100	33	11.6	57
GA	701,600	15	44,600	12	105,700	16	15.7	19
IL	316,600	30	15,700	20	45,500	26	20.1	37
IN	115,200	38	6,700	26	23,300	32	17.1	46
KY	305,400	53	14,200	23	47,300	49	21.5	58
LA	184,600	47	13,700	33	25,000	36	13.5	57
MD	65,100	22	5,200	24	14,600	22	12.5	32
MS	225,100	19	13,800	17	31,700	21	16.3	26
NC	662,300	30	46,800	20	113,700	24	14.1	36
OH	149,100	35	10,700	23	34,800	28	13.9	42
PA	142,900	37	14,200	29	50,100	35	10.1	47
RI	1,300	131	200	67	600	93	6.0	147
SC	555,200	37	29,900	22	90,400	31	18.6	43
TN	408,500	37	25,200	22	59,400	30	16.2	43
VA	208,600	35	15,300	23	40,700	33	13.7	42
WI	45,500	30	5,100	33	17,000	34	8.9	45
WV	14,900	77	1,100	27	3,100	39	14.0	82
Central	7,334,600	14	430,400	†	1,344,400	13	†	†
AR	258,200	29	16,300	28	36,200	27	15.9	41
CO	141,200	20	13,100	18	29,700	19	10.8	27
IA	128,100	19	9,700	15	25,300	17	13.2	24
KS	427,600	18	28,600	12	77,200	17	14.9	22
MN	96,700	79	6,500	58	18,000	55	15.0	98
MO	321,600	20	25,200	14	65,100	21	12.8	24
MT	16,000	53	1,900	44	3,500	43	8.6	69
NE	132,000	22	9,700	19	24,500	18	13.7	29
NM	47,900	26	4,400	18	12,800	33	10.8	31
ND	76,900	30	5,300	24	15,800	35	14.5	39
OK	400,400	28	23,800	14	58,500	21	16.8	32
SD	112,400	46	5,600	22	17,100	33	20.1	51
TX	5,155,300	19	278,700	13	956,800	18	18.5	23
WY	20,100	40	1,700	27	3,700	36	11.5	48
Western	1,561,400	10	104,300	†	297,000	10	†	†
AZ	395,800	12	21,900	6	62,800	9	18.1	13
CA	900,200	15	57,100	10	164,100	16	15.8	18
ID	108,900	33	7,300	29	25,300	38	15.0	44
NV	32,100	33	3,200	23	7,600	28	10.1	40
OR	27,200	40	4,300	35	9,000	31	6.3	53
UT	36,100	32	6,000	26	15,300	38	6.0	41
WA	61,100	30	4,600	23	12,800	30	13.4	38
United States	13,502,000	8	837,800	†	2,431,000	8	†	†

<sup>a</sup>Hunter number estimates at the management unit and national levels may be biased high, because the HIP sample frames are state specific; therefore hunters are counted more than once if they hunt in >1 state. Variance is inestimable.

<sup>b</sup>Seasonal harvest per hunter.

<sup>c</sup>No estimate available.

**Table 4.** Number of mourning doves banded in each management unit, state, and year, 2003–2016. Only known-age birds banded in July or August are included in the table and used in analysis of survival and harvest rates.

Mgmt Unit											
State	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013
Eastern	15,652	17,454	20,142	20,862	21,717	19,461	21,309	20,475	18,946	19,525	19,411
AL	1,130	1,112	991	961	889	117	1,147	1,026	942	1,010	1,097
DE	0	0	0	0	0	68	111	133	103	205	107
FL	830	960	916	858	773	1,027	799	865	736	968	805
GA	1,424	1,161	1,396	1,136	1,234	1,332	1,450	1,670	1,244	1,498	1,258
IL	6	6	47	1,163	1,267	1,378	1,877	1,833	2,034	1,501	1,276
IN	6	1,175	1,211	1,253	1,261	963	1,008	1,312	1,162	1,418	1,136
KY	1,444	1,566	1,454	1,637	1,608	1,867	2,391	2,232	1,786	1,299	1,553
LA	1,205	655	2,412	2,581	3,516	2,347	1,955	1,826	1,738	1,362	1,729
MD	472	482	719	571	708	322	334	312	377	346	366
MI	39	26	0	2	6	2	4	0	2	10	0
MS	1,071	994	1,008	656	690	822	928	448	462	605	666
North Atl. <sup>a</sup>	20	4	19	34	12	12	460	1,176	1,286	967	974
NC	1,283	1,539	1,662	1,299	1,307	1,736	1,685	1,198	795	1,847	1,734
OH	1,984	2,712	2,020	1,976	1,993	1,958	2,007	955	1,264	1,393	1,300
PA	1,564	1,590	1,658	1,838	1,748	942	903	899	827	899	1,007
RI	0	2	0	0	0	0	14	22	0	0	13
SC	1,041	863	1,484	1,461	1,761	1,720	1,875	1,953	1,911	1,795	1,902
TN	938	1,277	1,154	1,275	866	1,199	653	854	635	651	785
VA	474	546	804	585	642	603	599	554	496	522	420
WI	7	18	561	973	836	725	761	838	807	926	895
WV	714	768	626	603	600	321	348	369	339	303	388
Central	10,491	12,562	10,960	11,355	10,499	16,230	19,595	17,380	18,710	18,219	18,868
AR	782	975	1,085	914	822	711	514	0	424	222	297
CO	7	12	11	20	467	753	670	953	984	940	1,254
IA	1,940	2,191	2,458	1,099	987	1,694	1,238	1,078	2,216	2,089	1,649
KS	1,230	1,426	1,412	1,457	1,099	2,377	3,388	2,445	3,211	3,385	3,739
MN	0	4	0	0	363	529	700	1,164	853	1,026	1,390
MO	1,983	2,063	1,739	2,219	1,729	2,512	2,861	2,903	2,296	2,168	2,453
MT	0	0	0	0	0	0	0	322	270	296	223
NE	926	1,237	721	753	799	1,057	1,014	997	1,316	1,454	1,345
NM	3	11	14	4	0	463	1,059	625	114	717	829
ND	745	1,293	1,072	976	703	782	1,135	1,666	1,741	1,433	1,344
OK	391	447	528	715	826	1,513	2,746	1,520	1,661	1,488	1,182
SD	1,506	1,303	851	1,768	1,456	1,713	1,693	1,771	1,356	1,430	1,370
TX	978	1,600	1,069	1,430	1,237	2,078	2,575	1,936	2,268	1,502	1,702
WY	0	0	0	0	11	48	2	0	0	69	91
Western	3,261	3,658	4,494	4,559	6,495	6,253	9,059	9,348	7,552	8,634	8,961
AZ	1,653	1,574	1,582	2,436	2,562	2,544	3,831	3,599	3,818	3,362	3,718
CA	252	157	819	1,160	1,870	1,706	2,693	3,468	1,422	2,458	2,269
ID	440	854	837	730	615	594	466	453	355	677	511
NV	0	0	0	0	0	120	431	488	642	729	200
OR	0	0	0	0	0	173	245	219	243	319	734
UT	0	0	0	233	722	398	685	553	323	319	770
WA	916	1,073	1,256	0	726	718	708	568	749	770	759
United States	29,404	33,674	35,596	36,776	38,711	41,944	49,963	47,203	45,208	46,378	47,240

<sup>a</sup>Combined total for North Atlantic non-hunt states: CT, NH, ME, MA, NJ, NY, and VT.

**Table 4** (continued). Number of mourning doves banded in each management unit, state, and year, 2003–2016. Only known-age birds banded in July or August are included in the table and used in analysis of survival and harvest rates.

Mgmt Unit	2014	2015	2016
Eastern	17,993	18,448	16,772
AL	1,149	987	1,133
DE	202	38	94
FL	906	772	759
GA	954	1,336	1,152
IL	1,988	2,048	1,810
IN	1,237	977	653
KY	1,430	1,759	1,324
LA	1,066	1,769	1,596
MD	279	306	221
MI	0	0	0
MS	791	675	448
North Atl. <sup>a</sup>	141	118	159
NC	1,326	1,163	1,199
OH	1,336	1,312	1,316
PA	993	795	737
RI	0	55	0
SC	1,831	1,990	1,918
TN	677	611	540
VA	525	580	442
WI	789	800	887
WV	373	357	384
Central	21,545	19,516	19,982
AR	342	300	359
CO	1,335	1,011	1,419
IA	1,960	2,027	1,906
KS	3,233	3,332	2,868
MN	782	388	357
MO	2,997	1,966	1,983
MT	417	439	283
NE	1,505	1,357	1,718
NM	661	701	682
ND	1,675	1,620	1,647
OK	1,561	1,604	1,402
SD	1,872	2,052	2,329
TX	2,770	2,391	2,645
WY	435	328	384
Western	10,139	10,951	9,110
AZ	3,319	2,983	3,032
CA	3,510	4,535	3,293
ID	756	770	685
NV	600	401	498
OR	1,122	1,057	737
UT	349	282	59
WA	483	923	806
United States	49,677	48,915	45,864

<sup>a</sup>Combined total for North Atlantic non-hunt states: CT, NH, ME, MA, NJ, NY, and VT.

**Table 5.** Estimates of mean annual survival and harvest rate of mourning doves by management unit and state that banded doves, 2003–2016. Estimates by age-class: hatch-year (HY) and after-hatch-year (AHY). Standard errors are in parentheses.

Management Unit State	Annual Survival				Annual Harvest Rate			
	HY (SE)		AHY (SE)		HY (SE)		AHY (SE)	
Eastern	0.28	(0.01)	0.41	(0.01)	0.087	(0.001)	0.060	(0.001)
AL	0.27	(0.01)	0.41	(0.02)	0.099	(0.009)	0.068	(0.006)
DE-MD <sup>a</sup>	0.26	(0.02)	0.39	(0.02)	0.130	(0.009)	0.092	(0.010)
FL	0.27	(0.02)	0.41	(0.02)	0.040	(0.005)	0.033	(0.006)
GA	0.27	(0.01)	0.41	(0.01)	0.128	(0.006)	0.078	(0.007)
IL	0.27	(0.01)	0.40	(0.02)	0.070	(0.003)	0.050	(0.005)
IN	0.27	(0.01)	0.40	(0.01)	0.080	(0.009)	0.075	(0.006)
KY	0.28	(0.01)	0.42	(0.01)	0.062	(0.004)	0.052	(0.004)
LA	0.31	(0.01)	0.45	(0.01)	0.114	(0.006)	0.062	(0.006)
MS	0.25	(0.01)	0.38	(0.02)	0.149	(0.009)	0.087	(0.005)
North Atl <sup>b</sup>	0.48	(0.07)	0.63	(0.07)	0.004	(0.001)	0.003	(0.001)
NC	0.24	(0.01)	0.37	(0.01)	0.103	(0.009)	0.067	(0.004)
OH	0.25	(0.01)	0.38	(0.01)	0.054	(0.004)	0.045	(0.004)
PA	0.26	(0.02)	0.39	(0.02)	0.045	(0.007)	0.023	(0.004)
SC	0.29	(0.01)	0.43	(0.01)	0.094	(0.006)	0.060	(0.004)
TN	0.24	(0.01)	0.36	(0.02)	0.112	(0.004)	0.077	(0.004)
VA	0.30	(0.02)	0.44	(0.02)	0.039	(0.006)	0.041	(0.004)
WI	0.33	(0.02)	0.47	(0.02)	0.060	(0.005)	0.036	(0.004)
WV	0.37	(0.04)	0.52	(0.04)	0.021	(0.003)	0.014	(0.004)
Central	0.28	(0.01)	0.45	(0.01)	0.070	(0.001)	0.055	(0.001)
AR	0.24	(0.02)	0.38	(0.02)	0.085	(0.014)	0.065	(0.006)
CO	0.40	(0.03)	0.57	(0.03)	0.013	(0.002)	0.029	(0.004)
IA	0.32	(0.01)	0.48	(0.01)	0.038	(0.009)	0.027	(0.008)
KS	0.32	(0.01)	0.48	(0.01)	0.068	(0.006)	0.060	(0.004)
MN	0.38	(0.03)	0.54	(0.03)	0.030	(0.006)	0.018	(0.005)
MO	0.20	(0.01)	0.33	(0.01)	0.169	(0.010)	0.141	(0.007)
MT	0.41	(0.08)	0.58	(0.08)	0.015	(0.006)	0.016	(0.006)
ND	0.44	(0.02)	0.61	(0.02)	0.021	(0.002)	0.013	(0.002)
NE	0.31	(0.02)	0.47	(0.02)	0.034	(0.004)	0.035	(0.003)
NM	0.45	(0.07)	0.62	(0.06)	0.007	(0.002)	0.008	(0.002)
OK	0.27	(0.01)	0.42	(0.02)	0.087	(0.007)	0.065	(0.010)
SD	0.38	(0.02)	0.55	(0.01)	0.036	(0.004)	0.028	(0.003)
TX	0.33	(0.02)	0.49	(0.01)	0.054	(0.006)	0.041	(0.005)
WY	0.11	(0.08)	0.20	(0.13)	0.007	(0.007)	0.020	(0.002)
Western	0.30	(0.01)	0.43	(0.01)	0.044	(0.001)	0.037	(0.001)
AZ	0.30	(0.02)	0.43	(0.02)	0.025	(0.003)	0.018	(0.002)
CA	0.30	(0.02)	0.42	(0.01)	0.058	(0.007)	0.066	(0.008)
ID	0.30	(0.03)	0.43	(0.03)	0.027	(0.004)	0.019	(0.003)
NV	0.36	(0.04)	0.49	(0.04)	0.051	(0.010)	0.038	(0.006)
OR	0.33	(0.04)	0.47	(0.04)	0.036	(0.011)	0.030	(0.005)
UT	0.27	(0.04)	0.39	(0.04)	0.022	(0.005)	0.015	(0.005)
WA	0.30	(0.02)	0.43	(0.02)	0.053	(0.006)	0.041	(0.009)

<sup>a</sup>Data combined for Delaware and Maryland.

<sup>b</sup>Data combined for North Atlantic states: CT, NH, ME, MA, NJ, NY, RI, and VT.

**Table 6.** Estimated age ratios (juveniles per adult) by management unit and state based on the Parts Collection Survey, 2007–2016. Age ratios are corrected for unknown age wings and differential vulnerability. Sample size is the number of wings examined. Standard errors are in parentheses.

Management Unit	2007 <sup>a</sup>		2008		2009		2010		2011		2012	
State												
Eastern	1.73	(0.04)	1.42	(0.03)	1.35	(0.03)	1.30	(0.02)	1.83	(0.04)	1.81	(0.04)
AL	3.79	(2.69)	1.25	(0.17)	1.95	(0.29)	1.35	(0.10)	2.14	(0.19)	2.74	(0.27)
DE	1.15	(0.16)	1.88	(0.23)	0.89	(0.18)	1.60	(0.24)	3.21	(0.45)	1.47	(0.17)
GA	3.13	(0.40)	1.70	(0.24)	1.43	(0.18)	1.77	(0.20)	3.51	(0.48)	2.09	(0.18)
IL	1.85	(0.11)	1.21	(0.08)	1.47	(0.11)	1.29	(0.08)	1.51	(0.12)	2.50	(0.21)
IN	1.62	(0.07)	1.80	(0.15)	1.54	(0.11)	1.15	(0.06)	2.00	(0.12)	1.60	(0.12)
KY	1.68	(0.14)	1.18	(0.17)	1.58	(0.17)	1.77	(0.14)	1.65	(0.12)	1.69	(0.14)
LA	1.09	(0.13)	1.61	(0.25)	2.26	(0.31)	2.30	(0.26)	2.94	(0.58)	1.60	(0.25)
MD	2.07	(0.21)	1.52	(0.19)	1.24	(0.13)	1.39	(0.12)	1.45	(0.14)	1.93	(0.15)
MS	1.42	(0.14)	1.57	(0.16)	1.81	(0.17)	1.07	(0.07)	1.38	(0.13)	1.70	(0.24)
NC	1.80	(0.14)	1.67	(0.14)	1.40	(0.09)	1.04	(0.05)	1.73	(0.13)	1.45	(0.09)
OH	2.06	(0.19)	2.26	(0.29)	1.42	(0.16)	0.87	(0.07)	1.75	(0.15)	2.36	(0.29)
PA	1.35	(0.14)	1.03	(0.11)	0.93	(0.10)	1.03	(0.11)	1.91	(0.24)	1.62	(0.18)
RI <sup>b</sup>	----	----	----	----	----	----	----	----	----	----	----	----
SC	1.91	(0.12)	1.39	(0.09)	1.17	(0.08)	1.55	(0.09)	2.37	(0.16)	1.50	(0.10)
TN	1.82	(0.28)	1.34	(0.20)	1.13	(0.11)	1.51	(0.14)	2.13	(0.21)	3.25	(0.36)
VA	1.79	(0.11)	1.23	(0.07)	0.88	(0.07)	1.19	(0.06)	1.38	(0.08)	1.58	(0.08)
WI	1.00	(0.18)	1.58	(0.17)	1.24	(0.18)	2.04	(0.23)	1.27	(0.19)	2.04	(0.27)
WV	1.93	(0.24)	2.56	(0.58)	1.16	(0.19)	1.62	(0.25)	2.09	(0.32)	1.39	(0.22)
Central	1.04	(0.02)	0.95	(0.02)	0.84	(0.02)	0.99	(0.02)	1.13	(0.02)	1.50	(0.03)
AR	1.09	(0.10)	2.77	(0.35)	1.27	(0.11)	1.19	(0.10)	1.52	(0.14)	2.54	(0.27)
CO	1.12	(0.06)	1.09	(0.07)	0.83	(0.06)	1.43	(0.09)	1.37	(0.10)	1.12	(0.11)
IA	†	†	†	†	†	†	†	†	2.07	(0.59)	1.54	(0.16)
KS	1.32	(0.07)	0.99	(0.07)	0.89	(0.07)	1.11	(0.07)	1.10	(0.07)	1.46	(0.11)
MN	1.26	(0.90)	0.54	(0.33)	2.51	(0.72)	6.41	(3.83)	0.98	(0.10)	2.06	(0.18)
MO	1.62	(0.12)	0.93	(0.07)	0.94	(0.06)	1.21	(0.10)	1.58	(0.11)	1.96	(0.13)
MT	1.30	(0.16)	0.68	(0.09)	1.45	(0.23)	1.49	(0.17)	1.85	(0.26)	1.27	(0.16)
ND	1.07	(0.15)	0.92	(0.11)	1.39	(0.26)	0.65	(0.09)	0.99	(0.10)	1.56	(0.16)
NE	0.68	(0.04)	0.83	(0.06)	0.80	(0.09)	1.02	(0.07)	0.82	(0.05)	1.49	(0.11)
NM	0.55	(0.08)	0.35	(0.04)	0.48	(0.04)	0.59	(0.04)	0.71	(0.07)	0.68	(0.06)
OK	1.41	(0.17)	1.35	(0.10)	1.15	(0.07)	1.05	(0.06)	1.76	(0.14)	1.72	(0.16)
SD	1.07	(0.09)	0.89	(0.07)	1.08	(0.11)	1.05	(0.10)	1.18	(0.11)	1.73	(0.15)
TX	0.78	(0.05)	1.24	(0.07)	0.67	(0.04)	0.86	(0.04)	1.21	(0.05)	1.47	(0.07)
WY	1.32	(0.16)	0.90	(0.10)	0.75	(0.10)	1.68	(0.16)	1.51	(0.14)	1.05	(0.13)
Western	1.05	(0.03)	1.29	(0.04)	1.17	(0.04)	1.15	(0.03)	1.11	(0.03)	1.34	(0.04)
AZ	0.52	(0.03)	0.85	(0.04)	0.72	(0.04)	0.74	(0.04)	0.74	(0.04)	0.72	(0.05)
CA	1.22	(0.08)	1.45	(0.08)	1.23	(0.10)	1.15	(0.06)	1.15	(0.06)	1.35	(0.07)
ID	1.12	(0.10)	0.88	(0.17)	1.52	(0.16)	1.56	(0.18)	1.45	(0.25)	1.56	(0.15)
NV	1.13	(0.11)	1.09	(0.21)	0.97	(0.13)	0.96	(0.08)	1.14	(0.11)	1.28	(0.13)
OR	1.75	(0.29)	1.42	(0.60)	1.10	(0.18)	2.24	(0.28)	0.98	(0.16)	0.98	(0.13)
UT	1.19	(0.16)	0.73	(0.09)	0.69	(0.14)	0.79	(0.09)	1.17	(0.11)	1.36	(0.19)
WA	1.50	(0.10)	1.62	(0.12)	1.55	(0.15)	1.41	(0.12)	1.53	(0.13)	1.66	(0.15)

† Iowa did not have a hunting season until 2011.

<sup>a</sup> Standard errors for estimates only incorporate sampling error for the proportion of young in the sample and do not incorporate additional uncertainty from correction factors for unknown age wings and differential vulnerability.

<sup>b</sup> Insufficient data to estimate age ratio for RI in most years.

**Table 6** (continued). Estimated age ratios (juveniles per adult) by management unit and state based on the Parts Collection Survey, 2007–2016. Age ratios are corrected for unknown age wings and differential vulnerability. Sample size is the number of wings examined. Standard errors are in parentheses.

Management Unit State									2007-2016		
	2013 <sup>a</sup>		2014		2015		2016		Sample Size	Mean	SE
Eastern	1.33	(0.03)	1.42	(0.04)	1.31	(0.04)	1.31	(0.05)	77,929	1.49	(0.01)
AL	1.67	(0.18)	1.10	(0.10)	1.56	(0.17)	1.86	(0.26)	3,781	1.65	(0.06)
DE	1.97	(0.37)	1.30	(0.21)	0.42	(0.11)	0.96	(0.26)	1,839	1.48	(0.07)
GA	1.45	(0.11)	1.70	(0.16)	1.30	(0.12)	1.69	(0.16)	4,184	1.79	(0.06)
IL	1.36	(0.11)	1.48	(0.12)	1.15	(0.12)	0.93	(0.12)	7,158	1.46	(0.04)
IN	1.49	(0.12)	1.28	(0.12)	1.05	(0.09)	0.93	(0.13)	8,989	1.49	(0.03)
KY	1.23	(0.10)	1.41	(0.12)	1.18	(0.15)	1.29	(0.18)	5,000	1.52	(0.04)
LA	1.82	(0.29)	1.01	(0.76)	5.29	(2.89)	0.86	(0.26)	1,619	1.78	(0.09)
MD	1.64	(0.18)	1.78	(0.25)	1.69	(0.29)	2.76	(0.58)	3,642	1.64	(0.06)
MS	1.19	(0.12)	1.38	(0.15)	1.50	(0.18)	0.96	(0.18)	4,166	1.35	(0.04)
NC	1.12	(0.08)	1.01	(0.09)	0.97	(0.08)	0.83	(0.10)	7,748	1.28	(0.03)
OH	1.35	(0.15)	2.14	(0.22)	0.95	(0.10)	1.59	(0.26)	4,027	1.50	(0.05)
PA	1.27	(0.17)	1.30	(0.23)	1.57	(0.26)	1.04	(0.19)	2,737	1.17	(0.05)
RI <sup>b</sup>	----	----	0.76	(0.76)	----	----	0.67	(0.61)	19	2.00	(0.97)
SC	1.28	(0.12)	1.88	(0.18)	1.94	(0.23)	2.85	(0.35)	7,832	1.66	(0.04)
TN	1.38	(0.16)	2.01	(0.25)	1.36	(0.16)	1.19	(0.31)	3,114	1.67	(0.06)
VA	0.98	(0.09)	1.16	(0.15)	2.35	(0.31)	0.92	(0.11)	8,633	1.32	(0.03)
WI	1.64	(0.20)	1.39	(0.19)	2.78	(0.55)	3.14	(0.84)	2,148	1.58	(0.07)
WV	0.95	(0.32)	3.98	(1.19)	2.74	(0.71)	0.94	(0.23)	1,312	1.71	(0.10)
Central	1.16	(0.03)	1.12	(0.03)	0.99	(0.03)	1.07	(0.05)	66,067	1.07	(0.01)
AR	1.51	(0.15)	0.82	(0.10)	1.27	(0.15)	1.15	(0.17)	3,997	1.39	(0.04)
CO	1.62	(0.15)	1.48	(0.14)	0.92	(0.07)	1.11	(0.18)	7,158	1.17	(0.03)
IA	1.26	(0.21)	1.16	(0.13)	0.78	(0.09)	0.88	(0.19)	1,334	1.07	(0.06)
KS	1.37	(0.20)	1.50	(0.13)	1.00	(0.08)	1.00	(0.17)	7,053	1.14	(0.03)
MN	1.24	(0.16)	1.45	(0.25)	1.05	(0.21)	1.15	(0.41)	1,630	1.30	(0.07)
MO	1.07	(0.12)	1.93	(0.26)	2.41	(0.31)	1.17	(0.23)	5,714	1.36	(0.04)
MT	1.40	(0.26)	1.42	(0.26)	0.98	(0.12)	0.53	(0.14)	2,081	1.18	(0.05)
ND	1.23	(0.13)	1.24	(0.13)	1.32	(0.11)	1.00	(0.23)	2,987	1.13	(0.04)
NE	0.82	(0.08)	0.77	(0.10)	0.81	(0.09)	1.21	(0.23)	5,951	0.86	(0.02)
NM	0.52	(0.07)	0.41	(0.06)	0.77	(0.14)	0.84	(0.21)	3,778	0.56	(0.02)
OK	1.75	(0.19)	0.89	(0.10)	1.32	(0.15)	1.78	(0.29)	5,642	1.31	(0.04)
SD	1.07	(0.10)	0.93	(0.08)	0.91	(0.09)	0.97	(0.20)	4,603	1.07	(0.03)
TX	1.40	(0.11)	1.56	(0.10)	1.14	(0.10)	1.22	(0.16)	11,196	1.09	(0.02)
WY	2.06	(0.33)	0.89	(0.10)	0.81	(0.08)	2.27	(1.74)	2,943	1.14	(0.04)
Western	1.72	(0.08)	1.33	(0.06)	1.35	(0.05)	1.03	(0.06)	34,965	1.22	(0.01)
AZ	1.38	(0.13)	0.75	(0.05)	0.97	(0.06)	0.79	(0.06)	11,184	0.73	(0.01)
CA	1.62	(0.16)	1.54	(0.12)	1.41	(0.12)	1.44	(0.20)	9,468	1.29	(0.03)
ID	1.64	(0.17)	1.58	(0.17)	1.68	(0.21)	1.06	(0.15)	3,127	1.42	(0.05)
NV	1.30	(0.23)	0.93	(0.15)	1.57	(0.23)	0.58	(0.26)	2,650	1.11	(0.04)
OR	1.52	(0.18)	1.77	(0.39)	1.43	(0.26)	1.35	(0.34)	1,602	1.45	(0.07)
UT	1.27	(0.21)	1.70	(0.25)	0.85	(0.12)	0.76	(0.20)	2,168	1.01	(0.04)
WA	2.20	(0.26)	2.30	(0.48)	1.87	(0.25)	0.68	(0.16)	4,766	1.59	(0.05)

<sup>a</sup> Standard errors for estimates only incorporate sampling error for the proportion of young in the sample and do not incorporate additional uncertainty from correction factors for unknown age wings and differential vulnerability.

<sup>b</sup> Insufficient data to estimate age ratio for RI in most years.

**Table 7.** Estimates of absolute abundance of mourning doves on 1 September each year based on band recovery and harvest data by year and management unit in the U.S., 2003–2016.

Year	Management Unit						Total (United States)	
	Eastern		Central		Western		N	SE
	N	SE	N	SE	N	SE		
2003	95,410,137	5,943,665	113,292,709	8,806,487	130,553,174	23,645,800	339,256,021	25,923,064
2004	83,996,203	3,697,045	211,841,057	14,346,492	85,330,402	10,815,982	381,167,662	18,343,267
2005	132,883,623	5,530,276	191,696,765	14,037,207	38,373,428	3,850,362	362,953,816	15,570,883
2006	89,850,049	3,610,435	198,878,459	13,131,452	49,983,023	4,595,415	338,711,531	14,373,173
2007	102,380,934	4,595,082	157,491,832	10,098,337	59,860,570	4,387,999	319,733,336	11,930,873
2008	98,054,573	4,040,673	169,328,484	10,710,906	52,516,245	4,289,543	319,899,303	12,225,004
2009	103,089,071	4,237,048	148,487,151	8,868,563	50,903,066	3,438,976	302,479,288	10,412,999
2010	89,878,190	4,158,553	149,488,117	9,512,177	54,722,323	3,827,152	294,088,630	11,064,455
2011	85,742,115	4,454,969	125,454,975	6,963,865	51,056,398	3,866,139	262,253,488	9,126,291
2012	86,822,493	4,426,412	148,811,443	12,058,816	69,355,734	5,485,348	304,989,670	13,967,720
2013	85,541,482	5,400,755	124,308,636	8,245,304	48,016,677	3,620,680	257,866,796	10,500,596
2014	67,619,636	3,438,223	160,404,627	9,525,464	45,181,169	3,304,695	273,205,432	10,652,551
2015	63,349,787	3,292,117	164,796,552	9,439,559	36,392,969	2,456,931	264,539,309	10,294,650
2016	62,313,189	3,522,284	171,020,595	13,605,742	45,773,738	3,494,026	279,107,522	14,482,089

**Appendix A.** Federal framework dates, season length, and daily bag limit for mourning dove hunting in the U.S. by management unit, 1918–2017.

Year	Management Unit								
	Eastern			Central			Western		
	Dates <sup>a</sup>	Days	Bag	Dates	Days	Bag	Dates	Days	Bag
1918	Sep 1–Dec 31	107	25	Sep 1–Dec 15	106	25	Sep 1–Dec 15	106	25
1919–22	Sep 1–Jan 31	108	25	Sep 1–Dec 15	106	25	Sep 1–Dec 15	106	25
1923–28	Sep 1–Jan 31	108	25	Sep 1–Dec 31	106	25	Sep 1–Dec 15	106	25
1929	Sep 1–Jan 31	106	25	Sep 1–Dec 31	106	25	Sep 1–Dec 15	106	25
1930	Sep 1–Jan 31	108	25	Sep 1–Dec 15	106	25	Sep 1–Dec 15	106	25
1931	Sep 1–Jan 31	106	25	Sep 1–Dec 15	106	25	Sep 1–Dec 15	106	25
1932–33	Sep 1–Jan 31	106	18	Sep 1–Dec 15	106	18	Sep 1–Dec 15	106	18
1934	Sep 1–Jan 31	106	18	Sep 1–Jan 15	106	18	Sep 1–Dec 15	106	18
1935	Sep 1–Jan 31	107	20	Sep 1–Jan 16	106	20	Sep 1–Jan 05	107	20
1936	Sep 1–Jan 31	77	20	Sep 1–Jan 16	76	20	Sep 1–Nov 15	76	20
1937 <sup>b</sup>	Sep 1–Jan 31	77	15	Sep 1–Nov 15	76	15	Sep 1–Nov 15	76	15
1938	Sep 1–Jan 31	78	15	Sep 1–Nov 15	76	15	Sep 1–Nov 15	76	15
1939	Sep 1–Jan 31	78	15	Sep 1–Jan 31	77	15	Sep 1–Nov 15	76	15
1940	Sep 1–Jan 31	77	12	Sep 1–Jan 31	76	12	Sep 1–Nov 15	76	12
1941	Sep 1–Jan 31	62	12	Sep 1–Oct 27	42	12	Sep 1–Oct 12	42	12
1942	Sep 1–Oct 15	30	10	Sep 1–Oct 27	42	10	Sep 1–Oct 12	42	10
1943	Sep 1–Dec 24	30	10	Sep 1–Dec 19	42	10	Sep 1–Oct 12	42	10
1944	Sep 1–Jan 20	58	10	Sep 1–Jan 20	57	10	Sep 1–Oct 25	55	10
1945	Sep 1–Jan 31	60	10	Sep 1–Jan 31	60	10	Sep 1–Oct 30	60	10
1946	Sep 1–Jan 31	61	10	Sep 1–Jan 31	60	10	Sep 1–Oct 30	60	10
1947–48 <sup>c</sup>	Sep 1–Jan 31	60	10	Sep 1–Dec 3	60	10	Sep 1–Oct 30	60	10
1949	Sep 1–Jan 15	30	10	Sep 1–Nov 14	45	10	Sep 1–Oct 15	45	10
1950	Sep 1–Jan 15	30	10	Sep 1–Dec 3	45	10	Sep 1–Oct 15	45	10
1951	Sep 1–Jan 15	30	8	Sep 1–Dec 24	42	10	Sep 1–Oct 15	45	10
1952	Sep 1–Jan 10	30	8	Sep 1–Nov 6	42	10	Sep 1–Oct 12	42	10
1953	Sep 1–Jan 10	30	8	Sep 1–Nov 9	42	10	Sep 1–Oct 12	42	10
1954 <sup>d</sup>	Sep 1–Jan 10	40	8	Sep 1–Nov 9	40	10	Sep 1–Oct 31	40	10
1955	Sep 1–Jan 10	45	8	Sep 1–Nov 28	45	10	Sep 1–Dec 31	45	10
1956 <sup>e</sup>	Sep 1–Jan 10	55	8	Sep 1–Jan 10	55	10	Sep 1–Jan 10	50	10
1957	Sep 1–Jan 10	60	10	Sep 1–Jan 10	60	10	Sep 1–Jan 10	50	10
1958–59	Sep 1–Jan 15	65	10	Sep 1–Jan 15	65	10	Sep 1–Jan 15	50	10
1960–61 <sup>f</sup>	Sep 1–Jan 15	70 <sup>g</sup>	12	Sep 1–Jan 15	60	15	Sep 1–Jan 15	50	10
1962	Sep 1–Jan 15	70 <sup>g</sup>	12	Sep 1–Jan 15	60	12	Sep 1–Jan 15	50	10
1963	Sep 1–Jan 15	70 <sup>g</sup>	10	Sep 1–Jan 15	60	10	Sep 1–Jan 15	50	10
1964–67	Sep 1–Jan 15	70 <sup>g</sup>	12	Sep 1–Jan 15	60	12	Sep 1–Jan 15	50	12
1968	Sep 1–Jan 15	70 <sup>g</sup>	12	Sep 1–Jan 15	60	12	Sep 1–Jan 15	50	10
1969–70	Sep 1–Jan 15	70 <sup>g</sup>	18 <sup>h</sup>	Sep 1–Jan 15	60	10	Sep 1–Jan 15	50	10
1971–79	Sep 1–Jan 15	70 <sup>g</sup>	12	Sep 1–Jan 15	60	10	Sep 1–Jan 15	50	10
1980	Sep 1–Jan 15	70	12	Sep 1–Jan 15 <sup>i</sup>	60	10	Sep 1–Jan 15	70 <sup>j</sup>	10 <sup>k</sup>
1981	Sep 1–Jan 15	70	12	Sep 1–Jan 15 <sup>i</sup>	45 <sup>l</sup>	15 <sup>l</sup>	Sep 1–Jan 15	70 <sup>j</sup>	10 <sup>k</sup>
1982	Sep 1–Jan 15	45 <sup>m</sup>	15 <sup>m</sup>	Sep 1–Jan 15 <sup>i</sup>	45 <sup>m</sup>	15 <sup>m</sup>	Sep 1–Jan 15	45 <sup>m</sup>	15 <sup>m</sup>
1983–86	Sep 1–Jan 15	60 <sup>m</sup>	15 <sup>m</sup>	Sep 1–Jan 15 <sup>i</sup>	60 <sup>m</sup>	15 <sup>m</sup>	Sep 1–Jan 15	60 <sup>m</sup>	15 <sup>m</sup>
1987–07 <sup>n</sup>	Sep 1–Jan 15	60 <sup>m</sup>	15 <sup>m</sup>	Sep 1–Jan 15 <sup>i</sup>	60 <sup>m</sup>	15 <sup>m</sup>	Sep 1–Jan 15	60 <sup>o</sup>	10
2008	Sep 1–Jan 15	70	15	Sep 1–Jan 15 <sup>i</sup>	60 <sup>m</sup>	15 <sup>m</sup>	Sep 1–Jan 15	60 <sup>o</sup>	10
2009–13	Sep 1–Jan 15	70	15	Sep 1–Jan 15 <sup>i</sup>	70	15	Sep 1–Jan 15	60 <sup>o</sup>	10
2014	Sep 1–Jan 15	90	15	Sep 1–Jan 15 <sup>i</sup>	70	15	Sep 1–Jan 15	60 <sup>o</sup>	15
2015	Sep 1–Jan 15	90	15	Sep 1–Jan 15 <sup>i</sup>	70	15	Sep 1–Jan 15	60	15 <sup>p</sup>
2016–17	Sep 1–Jan 15	90	15	Sep 1–Jan 15 <sup>i</sup>	90	15	Sep 1–Jan 15	60	15 <sup>p</sup>

<sup>a</sup> From 1918–1947, seasons for doves and other “webless” species were selected independently and the dates were the earliest opening and latest closing dates chosen. Dates were inclusive. There were different season lengths in various states with some choosing many fewer days than others. Only bag and possession limits, and season dates were specified.

<sup>b</sup> Beginning in 1937, the bag and possession limit included white-winged doves in selected states.

<sup>c</sup> From 1948–1953, states permitting dove hunting were listed by waterfowl flyway. Only bag and possession limits, and season dates were specified.

<sup>d</sup> In 1954–1955, states permitting dove hunting were listed separately. Only bag and possession limits, and season dates were specified.

<sup>e</sup> From 1956–1959, states permitting dove hunting were listed separately. Framework opening and closing dates for seasons (but no maximum days for season length) were specified for the first time along with bag and possession limits.

**Appendix A. Continued.**

<sup>f</sup> In 1960, states were grouped by management unit for the first time. Maximum season length was specified for the first time.

<sup>g</sup> Half days.

<sup>h</sup> More liberal limits allowed in conjunction with an Eastern Management Unit hunting regulations experiment.

<sup>i</sup> The framework extended to January 25 in Texas.

<sup>j</sup> 50–70 days depending on state and season timing.

<sup>k</sup> Arizona was allowed 12.

<sup>l</sup> States had the option of a 60-day season and daily bag limit of 12.

<sup>m</sup> States had the option of a 70-day season and daily bag limit of 12.

<sup>n</sup> Beginning in 2002, the limits included white-winged doves in all states in the Central Management Unit. Beginning in 2006, the limits included white-winged doves in all states in the Eastern Management Unit.

<sup>o</sup> 30–60 days depending on state (30 in Idaho, Nevada, Oregon, Utah, Washington; 60 in Arizona and California).

<sup>p</sup> In Idaho, Nevada, Oregon, and Utah daily limit is 15 mourning and white-winged doves in the aggregate. In Arizona and California daily limit is 15 mourning and white-winged doves in the aggregate, of which no more than 10 can be white-winged doves.

**U.S. Fish and Wildlife Service  
Division of Migratory Bird Management  
Population and Habitat Assessment Branch  
11510 American Holly Drive  
Laurel, Maryland 20708-4002  
<http://www.fws.gov>**

**August. 2017**

**For StateTransfer Relay Service: TTY/  
Voice:711**