

**POST-CONSTRUCTION AVIAN AND BAT  
FATALITY MONITORING AND GRASSLAND  
BIRD DISPLACEMENT SURVEYS AT THE  
JUDITH GAP WIND ENERGY PROJECT,  
WHEATLAND COUNTY, MONTANA**

*Prepared for*

**Judith Gap Energy, LLC**  
Chicago, Illinois

*Prepared by*

**TRC Environmental Corporation**  
Laramie, Wyoming

January 2008



**POST-CONSTRUCTION AVIAN AND BAT FATALITY MONITORING  
AND GRASSLAND BIRD DISPLACEMENT SURVEYS  
AT THE JUDITH GAP WIND ENERGY PROJECT,  
WHEATLAND COUNTY, MONTANA**

Prepared for

**Judith Gap Energy, LLC  
Chicago, Illinois**

By

**TRC Environmental Corporation  
Laramie, Wyoming  
TRC Project 51883-01 (112416)**

**January 2008**

---

## EXECUTIVE SUMMARY

Judith Gap Energy, LLC operates the Judith Gap Wind Energy Project (JGWEP) in Wheatland County, Montana. The Farm Service Agency and the Montana State Lands Department were cooperating agencies for the preparation of an environmental assessment for the project. One of the stipulations for the Finding of No Significant Impact was a commitment to monitor avian and bat fatalities during operations. The *Avian and Bat Monitoring Plan, Judith Gap Wind Energy Project, Wheatland County, Montana* (TRC 2006) presented a detailed protocol for monitoring bird and bat fatalities at the JGWEP. Studies were conducted during the fall 2006 and spring 2007 migration periods, and this report presents the results and fatality estimates for birds and bats at JGWEP. A grassland bird displacement study was also conducted, and comparisons were made between point counts of breeding grassland birds before and after construction and operations of the JGWEP.

The JGWEP consists of 90 turbines, 20 of which were selected for monthly carcass searches. Estimates of total fatalities and fatalities per turbine were calculated based on the number of carcasses found after adjusting for searcher efficiency and scavenger removal rates. Thirty-five bat and 15 avian carcasses were found during carcass searches, and one bat and 11 avian carcasses were found incidentally. Searcher efficiency estimates were 17% for small birds (42% when adjusted for carcass placement-searcher efficiency trial time gap) and 77% for large birds. Estimated mean carcass residency time was 9.07 days for small birds and 15.43 days for large birds. Based on the number of carcasses found, and adjusting for searcher efficiency and scavenger removal rates, estimated turbine-related fatalities at JGWEP during the study period were 1,206 bats (13.40/turbine) and 406 birds (4.52/turbine). Bird fatality estimates were further broken down into small birds (344 fatalities; 3.83/turbine) and large birds (62 fatalities; 0.69/turbine). Bird fatalities were fairly uniformly spread over the study period, but 97% of the bat fatalities were recorded during fall migration (71% in August). One raptor fatality, a short-eared owl, was recorded during the study period. The results of the grassland bird displacement study suggested that construction of the windfarm did not negatively impact numbers of breeding grassland birds. Point counts performed in 2003 before construction of the windfarm were

replicated in 2007, and there was a 54% increase in numbers of birds detected in the vicinity of the turbines compared to a 20% increase in control plots with no turbines.

## **1.0 INTRODUCTION**

Judith Gap Energy, LLC (JGE) operates a 135-megawatt (MW) wind energy facility consisting of 90 turbines on approximately 8,000 acres (3,237 ha) in Wheatland County, Montana (Figure 1.1). The turbines are GE 1.5SLE 1.5-MW models with a 262-ft (80-m) hub height, 253-ft (77-m) blade diameter, and a rotor-swept area of 50,129 ft<sup>2</sup> (4,657 m<sup>2</sup>). The facility became operational in November 2005 and all turbines were on-line by January 2006. The Judith Gap Wind Energy Project (JGWEP) interconnects with NorthWestern Energy's transmission lines and provides as much as 8% of NorthWestern's electric supply.

Studies at existing wind farms in California have suggested that wind power development may detrimentally impact birds and bats due to mortality from collisions with turbine rotors and collisions with or electrocution by associated transmission lines (McCrary et al. 1986; Howell and DiDonato 1991; Orloff and Flannery 1992). Birds also may be negatively impacted by displacement resulting from habitat loss and disturbance (Leddy et al. 1999; Strickland 2004). In general, studies conducted at wind energy facilities outside California have found lower rates of avian and bat fatalities (Higgins et al. 1996; Erickson et al. 2000; Young et al. 2003; Erickson 2004); however, studies conducted at wind farms located on mountain ridgetops in the eastern U.S. have reported high fatality rates for migrating bats (Kerns and Kerlinger 2004; Fiedler et al. 2007). This suggests that local species composition and abundance, geographic area, topography, and turbine type and placement all contribute to the potential for avian and bat fatalities at individual wind farms.

Prior to construction of the JGWEP, the Farm Service Agency and the Montana State Lands Department prepared an environmental assessment (EA) of the proposed project. Given the JGWEP's proposed location, there was concern that the topography north of the project area--a gap between two mountain ranges--might funnel migratory birds and bats through the project area, thereby elevating the potential for avian and bat collisions with wind turbines during the migration period. Although preconstruction surveys suggested that mortality rates would not exceed those documented at other western wind farms, JGE committed to monitoring avian and




Figure 1.1 Project Location.

---

bat fatalities during its operation as part of the stipulations in the Decision Record and Finding of No Significant Impact for the EA. JGE representatives met with the U.S. Fish and Wildlife Service and NorthWestern Energy on April 11, 2006, and agreed to a monitoring plan that incorporated the primary concerns of the involved parties and focused on the as-built project, which included 30 fewer turbines than the original project evaluated in the EA. The monitoring plan, which involved conducting fatality searches during bird migration periods, was also agreed upon by a Technical Advisory Committee (TAC) comprised of agency, energy, and consultant personnel. The monitoring plan included estimating the number of avian and bat fatalities attributable to the project based on avian and bat fatalities documented during migration periods. Fatality estimates were calculated by adjusting for two potential sources of bias: 1) the amount of time carcasses remained on-site (scavenger removal bias) and 2) differences in detection abilities by observers (searcher efficiency rates) (Morrison 2002).

Migrant and resident birds may be killed by colliding with wind turbines and associated energy-related structures, and resident birds may be displaced from wind facilities by loss of habitat or disturbance in the immediate vicinity of the turbines during and after construction (Leddy et al. 1999; Strickland 2004). To evaluate the potential displacement of resident birds by the JGWEP, point-count surveys were conducted from April-June 2007 along transects located both within the wind farm and in control areas outside the wind farm. Results of these point counts were compared to preconstruction point counts that were conducted along the same transects in 2003, and the results are presented in Appendix A.

---

## 2.0 METHODS

### 2.1 PROJECT AREA LOCATION

The JGWEP straddles north/south-trending State Highway 191 between the towns of Judith Gap (located 6.2 mi [10.0 km] north of the site) and Harlowtown (located 7.5 mi [12.0 km] south of the site) in Wheatland County, Montana. Topographically, the project area lies less than 10.0 mi (16.0 km) south of a 10 to 15-mi (16 to 24-km) wide gap between the Big Snowy Mountains and the Little Belt Mountains. The project area and its surroundings consist of rolling hills, ridges, and flat benches dissected by broad valleys at an elevation of approximately 4,500-4,600 ft (1,372-1,402 m). Wind turbines are located in Sections 1, 4, 5, 6, 7, 8, 9, 12, 17, and 18, T9N, R16E, and Sections 29, 30, 31, 32, and 33, T10N, R16E. Vegetation in the project area consists of native short-grass prairie, located primarily on the west side of Highway 191; cultivated dryland agriculture, located primarily east of the highway; and Conservation Reserve Program (CRP) grassland, located primarily east of Highway 191 and north of east/west-trending Oxford Road (Figure 2.1). Sixteen of the 90 turbines are located in cropland, 51 are in CRP, and 23 are in short-grass prairie.

### 2.2 FATALITY STUDIES

#### 2.2.1 Standardized Carcass Searches

Monthly carcass searches were conducted in the project area during the fall (August-October 2006) and spring (February-May 2007) migration seasons to enable estimation of the number of avian and bat fatalities attributable to collision with the wind turbines. Searches were conducted on survey plots that were established at 20 of the 90 wind turbines. The number of turbine plots that were sampled was based on an examination of the estimated confidence intervals attained under different sampling scenarios. Assuming a fatality rate of 4.5 fatalities per turbine per year (U.S. Farm Service Agency and Montana Department of State Lands 2004), and with a desired confidence level of 90%, it was determined that a sample size of 20 turbines would enable fatality estimates with a margin of error of approximately  $\pm 74$  birds/bats per year (18%).




Figure 2.1 Turbine Locations and Habitat Association at the Judith Gap Wind Energy Project.

---

Six of the sampled turbines were located in native grassland habitat and 14 were in CRP. No turbine plots were located in cropland because walking transects in these areas would have damaged crops. Data collected during carcass searches at other wind energy sites have shown that nearly all birds and bats that strike turbines fall within 207 ft (63 m) of the tower (Erickson et al. 2000; Johnson et al. 2002; Young et al. 2003); therefore, plot sizes at JGWEP were conservatively set at 623 x 623 ft (190 x 190 m) to ensure that birds and bats that struck a turbine fell within the search plot. Square plots (rather than circular plots) were used to facilitate marking the plot boundaries while conducting the searches. The long axis of each plot (i.e., the diagonal axis connecting two nonadjacent corners) was oriented east/west so that the maximum distances searched coincided with the direction of the prevailing winds.

To locate avian and bat carcasses, trained searchers walked along unmarked linear transects spaced approximately 20-33 ft (6-10 m) apart across the entire survey plot. Spacing of transects varied depending on the season, topography, and type of vegetation because these factors affected visibility. Searchers typically searched for birds along both sides of each transect out to approximately 13-16 ft (4-5 m), and spent approximately two hours searching each of the 20 turbine plots.

Carcasses found on turbine plots were assumed to have been killed by striking turbines unless evidence indicated otherwise (e.g., if the carcass was a juvenile bird that was too young to fly). This protocol results in a conservative estimate of wind project fatalities, and has become standard methodology for wind energy and other collision-related mortality studies (e.g., Erickson et al. 2003; Young et al. 2003; Erickson et al. 2000) because of the high costs associated with obtaining accurate estimates of natural and reference mortality rates.

The location of each carcass was recorded using a global positioning system (GPS) unit. Carcasses (or carcass remains) were photographed as they were found, marked with spray paint so they would not be counted again if they were relocated during a future search, and left on site. Searchers filled out a Fatality/Injured Wildlife Data Form (Appendix C) for every carcass found. Recorded data included turbine/survey plot number; carcass species; sex and age, if they




Figure 2.2 Carcass Search Plot Layout.

---

could be determined; date and time the carcass was located; condition of the carcass; approximate time since death; and, if the animal was estimated to have been dead less than one week, weather conditions that occurred at or before the estimated time of death. When in-field identification of carcass species was not possible, identification was made later using the photographs. Distances of carcasses from the search tower were calculated later in a geographic information system (GIS) using the carcass GPS location and the Measure function in ArcMap (ArcGIS 9.2). The condition of the carcass was categorized as follows.

- **Intact:** An intact carcass that was not badly decomposed and showed little or no signs of being fed upon by a predator or scavenger.
- **Scavenged:** An entire carcass that showed signs of being fed upon by a predator or scavenger, or a dismembered carcass (carcass remains such as wings, skeletal remains, or legs).
- **A feather spot:** Ten or more feathers or two or more primaries (flight feathers) present, indicating predation or scavenging of the carcass.

U.S. Fish and Wildlife Service enforcement personnel were notified by e-mail when a migratory bird was located during a carcass search survey or incidentally in the project area. Bats were collected and frozen for DNA analysis at the request of the Montana Fish Wildlife and Parks Department.

### **2.2.2 Incidental Carcass Finds**

Fatalities found within search plots but outside designated search times and fatalities found outside designated search plots were treated as incidental discoveries and were not included in fatality estimates. Fatalities documented by operations and maintenance personnel and others not conducting the standardized carcass searches were photographed to facilitate future identification and recorded on the Fatality/Injured Wildlife Data Form (Appendix C). Recorded data included location of the carcass; date and time the carcass was located; carcass species, if determined; condition of the carcass; distance and bearing of the carcass from the nearest turbine or structure; approximate time since death; and, if the animal was estimated to have been dead less than one week, weather conditions that occurred at or before the estimated time of death.

---

### **2.2.3 Searcher Efficiency Trials**

The objective of the searcher efficiency trials was to evaluate the searchers' detection abilities. Searcher efficiency trials were conducted on the same survey plots as the carcass searches and provided an estimate of the proportion of actual avian and bat carcasses that were likely to be detected by searchers. Searcher efficiency trials were conducted to evaluate searcher efficiency by season, habitat, and carcass size. The searcher efficiency rates were then used to adjust the estimated avian and bat fatality rates for searcher bias.

Searcher efficiency was anticipated to be low in CRP habitat because of vegetation reducing carcass visibility. However, it was decided not to improve detection of carcasses by mowing CRP search areas because mowing would have increased the turbine footprint, reduced available cover for grassland birds, and possibly increased carcass scavenging rates.

Six searcher efficiency trials were conducted: one in each month that turbine plots were searched, except in February when weather conditions and safety issues with the turbines disrupted a portion of the monitoring work. Searchers did not know when trials were being conducted, nor on which survey plots searcher efficiency carcasses had been placed. During each of the six searcher efficiency trials, 10 bird carcasses (five large and five small) were placed on randomly selected survey plots. Large bird carcasses were represented by hen pheasants, and small bird and bat carcasses were represented by small quail. Both carcass types were purchased from a game farm. Because no naturally occurring pheasants or quail were observed on-site, the use of these species for searcher efficiency trials prevented feathers scattered by scavengers from being erroneously recorded as turbine-strike fatalities. Searcher efficiency trials conducted previously (Johnson et al. 2003; Young et al. 2003) suggested that searcher efficiency rates were similar for bats and small birds. Therefore, small birds were used to determine the searcher efficiency rates for both small birds and bats in this study.

Carcasses were placed in random locations on the selected plots. Prior to placement, each carcass was marked discreetly so that it could be identified as a searcher efficiency carcass rather than a turbine-strike carcass. Because of personnel limitations and travel distances to the wind

---

farm, turbine plots containing searcher efficiency carcasses sometimes were not surveyed until several days after the carcasses had been placed. Each searcher efficiency carcass that was found was recorded, and searcher efficiency rates for large and small carcasses, CRP and grasslands, and fall and spring migration periods were calculated.

#### **2.2.4 Carcass Removal Trials**

Carcass removal trials were conducted to determine how long carcasses typically remained on the search plots (i.e., the length of time that they were available for detection). Carcass removal included removal by predators, scavengers, or any other means. Estimates of carcass removal were factored into overall fatality estimates to account for removal bias (i.e., turbine-strike fatalities being removed from the search area before they could be located by searchers).

Carcass removal trials were conducted six times during the monitoring study--three times during the fall migration period (August, September, and October) and three times during the spring migration period (February, March, and May)--to accommodate variations in weather, vegetation conditions, and scavenger densities and activities. Plots that were designated as carcass-search plots were excluded from scavenger removal trials because the presence of additional carcasses may have led to increased scavenger activity on the carcass-search plots. Each carcass removal trial consisted of monitoring the fate of bird carcasses that were placed in the project area. Ten bird carcasses (five large and five small) were distributed within 623 x 623-ft (190 x 190-m) plots (the same size as those used in the carcass searches) around randomly selected turbines. Large bird carcasses were represented by pheasants, and small bird and bat carcasses were represented by small quail. Trials conducted with bats in other studies (Young et al. 2003; Erickson et al. 2004; Fiedler 2004) suggested that carcass removal rates were similar for bats and small birds. Therefore, small birds were used to determine carcass removal rates for both small birds and bats at JGWEP.

To avoid attracting additional scavengers to a particular plot and to maintain the independence of the data, only one carcass was placed in a plot. Carcasses were monitored every day for the first 4 days post-placement, and then on days 7, 10, 14, and 20 post-placement. The presence or

---

absence of carcasses or carcass remains was noted each time the carcasses were monitored. Carcasses were categorized as Present/Unscavenged, Present/Scavenged, or Absent. After the 20-day carcass removal trial, all remaining carcasses were removed from the study plots and were recorded as having been present 21 days.

### **2.2.5 Statistical Analyses for Determining Fatality Rates**

The estimated total number of wind turbine-related fatalities at the JGWEP was based on four components: 1) the number of carcasses found; 2) the number of turbines searched compared to total number of turbines at the facility; 3) the searcher efficiency rate expressed as the proportion of planted carcasses found by searchers; and 4) the scavenger removal rate, which was expressed as the mean length of time a carcass remained in the study area and was indicative of the length of time fatalities might be available for detection by searchers.

#### **2.2.5.1 Number of Carcasses Documented During Carcass Searches**

The average number of carcasses detected per turbine was:

$$\bar{c} = \frac{\sum_{i=1}^k c_i}{k}$$

where  $c_i$  was the number of carcasses detected at turbine  $i$  for the period of study and  $k$  was the number of turbines searched.

#### **2.2.5.2 Estimation of Searcher Efficiency**

Searcher efficiency was expressed as  $p$ , the estimated proportion of placed (trial) carcasses found by searchers. Searcher efficiency rates were estimated by major habitat type, carcass size, and season.

### 2.2.5.3 Estimation of Scavenger Removal Rates

Carcass removal rates were expressed as the mean length of time that a trial carcass remained in the study area ( $\bar{t}$ ) before it was removed and was calculated as:

$$\bar{t} = \frac{\sum_{i=1}^s t_i}{s - s_c}$$

where  $t_i$  = the length of time a trial carcass remained in the study area before it was removed,  $s$  = the number of carcasses used in the scavenging trials, and  $s_c$  = the number of trial carcasses still present at 20 days. This estimator was used due to the right-censoring of data caused by removal of trial carcasses at 20 days.

### 2.2.5.4 Estimation of the Total Number of Facility-related Fatalities

Estimation of total mortality for bats and birds (large, small, and combined) at JGWEP involved an extrapolation of the number of carcasses found during searches, and incorporated other factors such as searcher efficiency and scavenger removal rates. The formula used to calculate total turbine-related fatalities ( $m$ ) was:

$$m = \left( \frac{N * I * C}{k * t * p} \right) \left( \frac{e^{I/t} - 1 + p}{e^{I/t} - 1} \right)$$

where  $N$  = the total number of turbines at JGWEP,  $I$  = the interval time between searches in days,  $C$  = total number of carcasses found,  $k$  = the number of turbines searched,  $t$  = the mean carcass residency time in days (i.e., mean number of days that carcasses remained on scavenger removal plots), and  $p$  = the searcher efficiency for the category (birds/bats) being calculated.

Ninety percent confidence intervals and a standard error of the fatality estimate were calculated using bootstrapping, a computer program that simulates repeated sampling (with replacement) of the collected data. A total of 5,000 iterations were used, and the confidence intervals were based on the 5th and 95th percentiles of the 5,000 bootstrap iterations. The standard deviation of the 5,000 iterations was used as the standard error of the estimated mortality.

---

### 3.0 RESULTS

A total of 62 carcasses was found at the JGWEP between July 2006-May 2007 (Figure 3.1; Table 3.1), including 36 (58% of total carcasses) bats and 26 (42%) birds. Fifty (81%) carcasses were found during carcass searches and 12 (19%) were found incidentally. Thirty-nine (78%) of the 50 carcasses found during searches were  $\leq 223$  ft (68 m) from the search turbine, three (6%) were 236-282 ft (72-86 m) from the turbine, and eight (16%) were  $\geq 318$  ft (97 m) from the turbine. The average distance of a carcass from the search turbine was 154 ft (47 m), with a range of 13 to 489 ft (4 to 149 m).

#### 3.1 CARCASS SEARCH SURVEYS

##### 3.1.1 Avian Carcass Finds

Fifteen avian carcasses, comprising at least seven species, were found in the 20 turbine plots that were searched monthly from August 2006 through October 2006 and February 2007 through May 2007--an average of 0.75 birds per turbine (Table 3.2). This represents 30% (15 of 50) of the total number of carcasses found on the study area during carcass searches, with the other 70% being bat carcasses. Avian carcasses were located an average of 210 ft (64 m) away from turbines, with a range of 56 to 489 ft (17 to 149 m) (Figure 3.1). Avian fatalities were documented at 11 (55%) of the 20 turbines that were searched monthly (Table 3.2). Two avian carcasses apiece were found at turbines 2, 32, 73, and 83, and one carcass was found at the other seven towers at which avian carcasses were found. No avian carcasses were found at turbines 6, 30, 40, 47, 53, 63, 66, 84, and 87.

Ten small birds (passerines) and five large birds (waterfowl and raptors) were found. Horned lark (four birds, 27% of all birds found) was the most common avian species located, followed by eared grebe and Wilson's warbler (two of each species, 13% of all birds found) (Table 3.1; Appendix B, Table B.1). One raptor, a short-eared owl carcass, was found. More bird carcasses (four) were found during August and September than in any other months (Figure 3.2). Avian carcasses were found in every month except February. A mean of 2.5 birds per month was found




Figure 3.1 Distribution of Carcasses Found During Carcass Searches and Incidentally at the Judith Gap Wind Energy Project.

Table 3.1 Number and Species of Bird and Bat Carcasses Found During Carcass Searches and Incidentally at the Judith Gap Wind Energy Project, July 2006-May 2007.

| Common Name            | Scientific Name                  | Number of Carcasses |
|------------------------|----------------------------------|---------------------|
| <b>Birds</b>           |                                  |                     |
| Eared grebe            | <i>Podiceps nigricollis</i>      | 5 <sup>1</sup>      |
| Western grebe          | <i>Aechmophorus occidentalis</i> | 2 <sup>2</sup>      |
| Mallard                | <i>Anas platyrhynchos</i>        | 1                   |
| Unidentified waterfowl | --                               | 1 <sup>3</sup>      |
| Merlin                 | <i>Falco columbarius</i>         | 1                   |
| Sharp-tailed grouse    | <i>Tympanuchus phasianellus</i>  | 1                   |
| American coot          | <i>Fulica americana</i>          | 4 <sup>4</sup>      |
| Short-eared owl        | <i>Asio flammeus</i>             | 1                   |
| Horned lark            | <i>Eremophila alpestris</i>      | 4 <sup>5</sup>      |
| Wilson's warbler       | <i>Wilsonia pusilla</i>          | 2                   |
| White-crowned sparrow  | <i>Zonotrichia albicollis</i>    | 1                   |
| Western meadowlark     | <i>Sturnella neglecta</i>        | 1                   |
| Unidentified passerine | --                               | 2                   |
| Subtotal Birds         |                                  | 26                  |
| <b>Bats</b>            |                                  |                     |
| Hoary bat              | <i>Lasiurus cinereus</i>         | 17 <sup>6</sup>     |
| Silver-haired bat      | <i>Lasionycteris noctivagans</i> | 4 <sup>7</sup>      |
| Unidentified bat       | --                               | 15                  |
| Subtotal Bats          |                                  | 36                  |
| Total Carcasses        |                                  | 62                  |

<sup>1</sup> Identification of one eared grebe was tentative. Alternatively, the bird may have been a horned grebe.

<sup>2</sup> Identification of western grebes was tentative. Alternatively, the birds may have been Clark's grebes.

<sup>3</sup> Tentative identification based on feather remains.

<sup>4</sup> Identification of one American coot was tentative and was based on minimal feather remains.

<sup>5</sup> Identification of two horned larks was tentative and was based on photographs of the scavenged carcass.

<sup>6</sup> Identification of one hoary bat was tentative and was based on photographs of pelage remnants.

<sup>7</sup> Identification of three silver-haired bats was tentative and was based on photographs of carcasses.

Table 3.2 Number and Type of Species Found at Turbine Plots During Carcass Searches at the Judith Gap Wind Energy Project, August-October 2006 and February-May 2007.<sup>1</sup>

| Species                | Turbine Number |   |   |    |    |    |    |    |    |    |    |    |    | Total Number of Carcasses |    |    |    |    |    |                |    |
|------------------------|----------------|---|---|----|----|----|----|----|----|----|----|----|----|---------------------------|----|----|----|----|----|----------------|----|
|                        | 1              | 2 | 6 | 30 | 31 | 32 | 34 | 40 | 47 | 50 | 53 | 63 | 64 |                           | 66 | 73 | 76 | 83 | 84 | 87             | 99 |
| <b>Birds</b>           |                |   |   |    |    |    |    |    |    |    |    |    |    |                           |    |    |    |    |    |                |    |
| Eared grebe            | 0              | 0 | 0 | 0  | 1  | 1  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0                         | 0  | 0  | 0  | 0  | 0  | 0              | 2  |
| Unidentified waterfowl | 0              | 0 | 0 | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 1  | 0                         | 0  | 0  | 0  | 0  | 0  | 0              | 1  |
| American coot          | 0              | 0 | 0 | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0                         | 0  | 0  | 0  | 0  | 0  | 1 <sup>2</sup> | 1  |
| Short-eared owl        | 0              | 1 | 0 | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0                         | 0  | 0  | 0  | 0  | 0  | 0              | 1  |
| Horned lark            | 0              | 1 | 0 | 0  | 0  | 0  | 1  | 0  | 0  | 0  | 0  | 0  | 0  | 0                         | 0  | 0  | 2  | 0  | 0  | 0              | 4  |
| Wilson's warbler       | 0              | 0 | 0 | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0                         | 2  | 0  | 0  | 0  | 0  | 0              | 2  |
| White-crowned sparrow  | 1              | 0 | 0 | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0                         | 0  | 0  | 0  | 0  | 0  | 0              | 1  |
| Western meadowlark     | 0              | 0 | 0 | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0                         | 0  | 1  | 0  | 0  | 0  | 0              | 1  |
| Unidentified passerine | 0              | 0 | 0 | 0  | 0  | 1  | 0  | 0  | 0  | 1  | 0  | 0  | 0  | 0                         | 0  | 0  | 0  | 0  | 0  | 0              | 2  |
| -----                  |                |   |   |    |    |    |    |    |    |    |    |    |    |                           |    |    |    |    |    |                |    |
| Subtotal Large Birds   | 0              | 1 | 0 | 0  | 1  | 1  | 0  | 0  | 0  | 0  | 0  | 0  | 1  | 0                         | 0  | 0  | 0  | 0  | 0  | 1              | 5  |
| -----                  |                |   |   |    |    |    |    |    |    |    |    |    |    |                           |    |    |    |    |    |                |    |
| Subtotal Small Birds   | 1              | 1 | 0 | 0  | 1  | 1  | 0  | 0  | 1  | 0  | 0  | 0  | 0  | 0                         | 2  | 1  | 2  | 0  | 0  | 0              | 10 |
| -----                  |                |   |   |    |    |    |    |    |    |    |    |    |    |                           |    |    |    |    |    |                |    |
| Subtotal Birds         | 1              | 2 | 0 | 0  | 1  | 2  | 1  | 0  | 0  | 1  | 0  | 0  | 1  | 0                         | 2  | 1  | 2  | 0  | 0  | 1              | 15 |
| <b>Bats</b>            |                |   |   |    |    |    |    |    |    |    |    |    |    |                           |    |    |    |    |    |                |    |
| Hoary bat              | 3              | 2 | 2 | 0  | 0  | 1  | 0  | 1  | 0  | 1  | 0  | 0  | 2  | 0                         | 2  | 0  | 1  | 2  | 0  | 0              | 17 |
| Silver-haired bat      | 0              | 0 | 0 | 0  | 0  | 0  | 1  | 0  | 0  | 0  | 0  | 1  | 0  | 1                         | 0  | 1  | 0  | 0  | 0  | 0              | 4  |
| Unidentified bat       | 0              | 0 | 2 | 0  | 1  | 0  | 0  | 2  | 0  | 0  | 0  | 1  | 1  | 2                         | 2  | 0  | 1  | 0  | 1  | 1              | 14 |
| -----                  |                |   |   |    |    |    |    |    |    |    |    |    |    |                           |    |    |    |    |    |                |    |
| Subtotal Bats          | 3              | 2 | 4 | 0  | 1  | 1  | 1  | 3  | 0  | 1  | 0  | 2  | 3  | 3                         | 4  | 1  | 2  | 2  | 1  | 1              | 35 |
| -----                  |                |   |   |    |    |    |    |    |    |    |    |    |    |                           |    |    |    |    |    |                |    |
| Total Carcasses        | 4              | 5 | 4 | 0  | 2  | 3  | 2  | 3  | 0  | 1  | 0  | 2  | 4  | 3                         | 6  | 2  | 4  | 2  | 1  | 2              | 50 |

<sup>1</sup> All turbine plots were searched seven times each (once a month) except for turbine 6, which was searched five times, and turbines 30, 31, 32, 34, 40, and 64, which were searched six times each. A total of 132 turbine-plot searches were conducted.

<sup>2</sup> An additional two American coots were found incidentally on the access road near turbine 99.




Figure 3.2 Number of Bird and Bat Carcasses Found by Month During Carcass Searches, August 2006-October 2006 and February 2007-May 2007.

during carcass surveys in the other six survey months excluding February. Carcass surveys were disrupted by weather and turbine safety issues during February; therefore, birds may have struck turbines during February but were not found. Mortality was higher in the fall (September and October) with an average of 3.0 avian carcasses found per month compared to an average of 1.7 avian carcasses found per month during spring (or an average of 1.3 carcass per month in the spring if February was included).

### **3.1.2 Bat Carcass Finds**

Thirty-five bat carcasses, comprising at least two species, were found in the 20 turbine plots that were searched between August 2006-October 2006 and February-May 2007 (Table 3.2). This represents 70% (35 of 50) of the carcasses found in the study area during carcass searches, or 56% (35 of 62) of the total number of carcasses found in the project area from July 2006-May 2007. Hoary bat (17 carcasses, 49% of bat carcasses) was the most common species located, followed by unidentified bat (14, 40%) and silver-haired bat (4, 11%). A mean of 1.75 bats was

---

found per turbine during the carcass surveys. Bat carcasses were located an average of 131 ft (40 m) away from turbines, with a range of 13 to 354 ft (4 to 108 m). Bat fatalities were documented at 17 (85%) of the 20 turbines that were searched monthly. Four bat carcasses each were found at turbines 6 and 73, and three bat carcasses each were found at turbines 1, 40, 64, and 66. No bat carcasses were found under turbines 30, 47, and 53, nor were any avian fatalities recorded at these three towers (see Table 3.2).

Substantially more bat carcasses were found in August (25, 71%) than in any other month, and more bat fatalities were documented in the fall (34, 97%) than in the spring (1, 3%) (see Figure 3.2).

## **3.2 INCIDENTAL FINDS**

### **3.2.1 Incidental Avian Finds**

Eleven avian carcasses (18% of the 62 bird and bat carcasses and 42% of the 26 bird carcasses found on the study area) were found incidentally (Table 3.3; Appendix B, Table B.1). All of the incidental bird finds were large birds (waterfowl, gallinaceous birds, and raptors), suggesting either that larger birds were more likely to be found or that scavengers removed smaller birds from the project area more quickly.

Two of the incidental avian carcasses (both western grebes) were found in July 2006 prior to initiation of the monitoring study (Appendix B, Table B.1). One additional carcass (an American coot) was found outside the monitoring period in early November 2006. Eight of the 11 incidental bird carcasses (73%) were found in the fall, and the three remaining bird carcasses found incidentally were found in May 2006 (Appendix B, Table B.1). An average of 1.6 birds per month was found incidentally during the summer/fall migration period compared to an average of 0.75 birds per month found incidentally during the winter/spring migration period.

When combining the 11 incidental avian finds with the 15 bird carcasses found during carcass searches, more eared grebes (five birds, 19% of the total number found) were found than any

Table 3.3 Number and Species of Carcasses Found Incidentally near Turbines and at Other Locations at the Judith Gap Wind Energy Project, July 2006 to May 2007.

| Common Name         | Turbine Numbers and Other Locations |    |    |    |    |    |    |                             | Total Number of Carcasses |                           |
|---------------------|-------------------------------------|----|----|----|----|----|----|-----------------------------|---------------------------|---------------------------|
|                     | 13                                  | 18 | 22 | 32 | 42 | 75 | 89 | Access Road near Turbine 99 |                           | O&M Building <sup>1</sup> |
| <b>Birds</b>        |                                     |    |    |    |    |    |    |                             |                           |                           |
| Eared grebe         | 2                                   | 0  | 0  | 0  | 0  | 0  | 1  | 0                           | 0                         | 3                         |
| Western grebe       | 0                                   | 2  | 0  | 0  | 0  | 0  | 0  | 0                           | 0                         | 2                         |
| Mallard             | 0                                   | 0  | 0  | 1  | 0  | 0  | 0  | 0                           | 0                         | 1                         |
| Merlin              | 0                                   | 0  | 0  | 0  | 0  | 0  | 0  | 0                           | 1                         | 1                         |
| Sharp-tailed grouse | 0                                   | 0  | 0  | 0  | 0  | 1  | 0  | 0                           | 0                         | 1                         |
| American coot       | 0                                   | 0  | 1  | 0  | 0  | 0  | 0  | 2                           | 0                         | 3                         |
| Subtotal Birds      | 2                                   | 2  | 1  | 1  | 0  | 1  | 1  | 2                           | 1                         | 11                        |
| <b>Bats</b>         |                                     |    |    |    |    |    |    |                             |                           |                           |
| Unidentified bat    | 0                                   | 0  | 0  | 0  | 1  | 0  | 0  | 0                           | 0                         | 1                         |
| Subtotal Bats       | 0                                   | 0  | 0  | 0  | 1  | 0  | 0  | 0                           | 0                         | 1                         |
| Total Carcasses     | 2                                   | 2  | 1  | 1  | 1  | 1  | 1  | 2                           | 1                         | 12                        |

<sup>1</sup> Carcass was found behind the Judith Gap Wind Energy facility Operations and Management (O&M) building.

other species, followed by American coots (4, 15%) and horned larks (4, 15%). More avian carcasses (nine carcasses) were found in September than in any other month (Figure 3.3).

### **3.2.2 Incidental Bat Finds**

The single bat carcass found incidentally during the course of the monitoring study was found in October 2006 (see Table 3.3; Appendix B, Table B.2). The remaining 35 (97%) bat carcasses found during the study were found during carcass searches.

## **3.3 SEARCHER EFFICIENCY BIAS**

Sixty bird carcasses (30 large birds and 30 small birds) were used for searcher efficiency trials during the study period. The interval between trial carcass placement and search dates averaged 8.3 days for small bird carcasses and 7.2 days for large bird carcasses. Searcher efficiency was




Figure 3.3 Number of Bird and Bat Carcasses Found by Month During Carcass Searches and Incidentally, August 2006-October 2006 and February 2007-May 2007.

77% for large bird carcasses and 17% for the small bird carcasses (Table 3.4). Results were similar across seasons except that large bird carcasses had a lower searcher efficiency during spring (67%) compared to summer/fall (87%). Large bird carcass searcher efficiency was similar in CRP (77%) and grassland (75%), but efficiency was higher for small bird carcasses in grassland (27%) compared to CRP (11%) (Table 3.5).

### 3.4 CARCASS REMOVAL BIAS

Sixty bird carcasses (30 large birds and 30 small birds) were used for scavenger removal trials during the study period (Tables 3.6 and 3.7). The average length of time carcasses remained was 12 days (15.4 days for large bird carcasses and 9.1 days for small bird carcasses). Seventeen percent of trial carcasses remained on the 20th day (23% of large bird carcasses and 10% of small bird carcasses). Mean carcass residency was shorter during the winter/spring period for both large bird (14.7 days) and small bird (5.3 days) carcasses when compared to summer/fall (16.3 days for large bird carcasses and 13.2 days for small bird carcasses) (see Table 3.6).

Table 3.4 Seasonal Searcher Efficiency Rates at Judith Gap Wind Energy Center.

| Searcher Efficiency Trial Date | Habitat <sup>1</sup> | Number of Large Birds Placed | Number of Large Birds Found | Large Bird Searcher Efficiency | Number of Small Birds Placed | Number of Small Birds Found | Small Bird Searcher Efficiency | Total Birds Placed | Total Found | Total Searcher Efficiency |
|--------------------------------|----------------------|------------------------------|-----------------------------|--------------------------------|------------------------------|-----------------------------|--------------------------------|--------------------|-------------|---------------------------|
| August                         | Grassland            | 1                            | 1                           | 100%                           | 2                            | 2                           | 100%                           | 3                  | 3           | 100%                      |
| August                         | CRP                  | 4                            | 3                           | 75%                            | 3                            | 0                           | 0%                             | 7                  | 3           | 43%                       |
| Total                          |                      | 5                            | 4                           | 80%                            | 5                            | 2                           | 40%                            | 10                 | 6           | 60%                       |
| September                      | Grassland            | 2                            | 2                           | 100%                           | 1                            | 0                           | 0%                             | 3                  | 2           | 67%                       |
| September                      | CRP                  | 3                            | 2                           | 67%                            | 4                            | 0                           | 0%                             | 7                  | 2           | 29%                       |
| Total                          |                      | 5                            | 4                           | 80%                            | 5                            | 0                           | 0%                             | 10                 | 4           | 40%                       |
| October                        | Grassland            | 1                            | 1                           | 100%                           | 2                            | 0                           | 0%                             | 3                  | 1           | 33%                       |
| October                        | CRP                  | 4                            | 4                           | 100%                           | 3                            | 1                           | 33%                            | 7                  | 5           | 71%                       |
| Total                          |                      | 5                            | 5                           | 100%                           | 5                            | 1                           | 20%                            | 10                 | 6           | 60%                       |
| Subtotal Summer/Fall           |                      | 15                           | 13                          | 87%                            | 15                           | 3                           | 20%                            | 30                 | 16          | 53%                       |
| March                          | Grassland            | 1                            | 1                           | 100%                           | 2                            | 0                           | 0%                             | 3                  | 1           | 33%                       |
| March                          | CRP                  | 4                            | 4                           | 100%                           | 3                            | 0                           | 0%                             | 7                  | 4           | 57%                       |
| Total                          |                      | 5                            | 5                           | 100%                           | 5                            | 0                           | 0%                             | 10                 | 5           | 50%                       |
| April                          | Grassland            | 1                            | 0                           | 0%                             | 2                            | 0                           | 0%                             | 3                  | 0           | 0%                        |
| April                          | CRP                  | 4                            | 1                           | 25%                            | 3                            | 0                           | 0%                             | 7                  | 1           | 14%                       |
| Total                          |                      | 5                            | 1                           | 20%                            | 5                            | 0                           | 0%                             | 10                 | 1           | 10%                       |
| May                            | Grassland            | 2                            | 1                           | 50%                            | 2                            | 1                           | 50%                            | 4                  | 2           | 50%                       |
| May                            | CRP                  | 3                            | 3                           | 100%                           | 3                            | 1                           | 33%                            | 6                  | 4           | 67%                       |
| Total                          |                      | 5                            | 4                           | 80%                            | 5                            | 2                           | 40%                            | 10                 | 6           | 60%                       |
| Subtotal Spring                |                      | 15                           | 10                          | 67%                            | 15                           | 2                           | 13%                            | 30                 | 12          | 40%                       |
| Total                          |                      | 30                           | 23                          | 77%                            | 30                           | 5                           | 17%                            | 60                 | 28          | 47%                       |

<sup>1</sup> CRP = Conservation Reserve Program land.

Table 3.5 Searcher Efficiency Rates in Different Habitats at Judith Gap Wind Energy Center.

| Habitat                | Searcher Efficiency Trial Date | Number of Large Birds Placed | Number of Large Birds Found | Large Bird Searcher Efficiency | Number of Small Birds Placed | Number of Small Birds Found | Small Bird Searcher Efficiency | Total Birds Placed | Total Birds Found | Total Searcher Efficiency |
|------------------------|--------------------------------|------------------------------|-----------------------------|--------------------------------|------------------------------|-----------------------------|--------------------------------|--------------------|-------------------|---------------------------|
| <b>CRP<sup>1</sup></b> |                                |                              |                             |                                |                              |                             |                                |                    |                   |                           |
|                        | August                         | 4                            | 3                           | 75%                            | 3                            | 0                           | 0%                             | 7                  | 3                 | 43%                       |
|                        | September                      | 3                            | 2                           | 67%                            | 4                            | 0                           | 0%                             | 7                  | 2                 | 29%                       |
|                        | October                        | 4                            | 4                           | 100%                           | 3                            | 1                           | 33%                            | 7                  | 5                 | 71%                       |
|                        | Subtotal Summer/Fall           | 11                           | 9                           | 82%                            | 10                           | 1                           | 10%                            | 21                 | 10                | 48%                       |
|                        | March                          | 4                            | 4                           | 100%                           | 3                            | 0                           | 0%                             | 7                  | 4                 | 57%                       |
|                        | April                          | 4                            | 1                           | 25%                            | 3                            | 0                           | 0%                             | 7                  | 1                 | 14%                       |
|                        | May                            | 3                            | 3                           | 100%                           | 3                            | 1                           | 33%                            | 6                  | 4                 | 67%                       |
|                        | Subtotal Spring                | 11                           | 8                           | 73%                            | 9                            | 1                           | 11%                            | 20                 | 9                 | 45%                       |
|                        | Total CRP                      | 22                           | 17                          | 77%                            | 19                           | 2                           | 11%                            | 41                 | 19                | 46%                       |
| <b>Grassland</b>       |                                |                              |                             |                                |                              |                             |                                |                    |                   |                           |
|                        | August                         | 1                            | 1                           | 100%                           | 2                            | 2                           | 100%                           | 3                  | 3                 | 100%                      |
|                        | September                      | 2                            | 2                           | 100%                           | 1                            | 0                           | 0%                             | 3                  | 2                 | 67%                       |
|                        | October                        | 1                            | 1                           | 100%                           | 2                            | 0                           | 0%                             | 3                  | 1                 | 33%                       |
|                        | Subtotal Summer/Fall           | 4                            | 4                           | 100%                           | 5                            | 2                           | 40%                            | 9                  | 6                 | 67%                       |
|                        | March                          | 1                            | 1                           | 100%                           | 2                            | 0                           | 0%                             | 3                  | 1                 | 33%                       |
|                        | April                          | 1                            | 0                           | 0%                             | 2                            | 0                           | 0%                             | 3                  | 0                 | 0%                        |
|                        | May                            | 2                            | 1                           | 50%                            | 2                            | 1                           | 50%                            | 4                  | 2                 | 50%                       |
|                        | Subtotal Spring                | 4                            | 2                           | 50%                            | 6                            | 1                           | 17%                            | 10                 | 3                 | 30%                       |
|                        | Total Grassland                | 8                            | 6                           | 75%                            | 11                           | 3                           | 27%                            | 19                 | 9                 | 47%                       |
|                        | Both Habitats Combined         | 30                           | 23                          | 77%                            | 30                           | 5                           | 17%                            | 60                 | 28                | 47%                       |

<sup>1</sup> CRP = Conservation Reserve Program land.

Table 3.6 Seasonal Scavenger Removal Rates at Judith Gap Wind Energy Project.

| Scavenger Efficiency Trial Date           | Habitat <sup>1</sup> | Percent of Carcasses Remaining After 20 Days | Mean Number of Days Carcasses Remained <sup>2,3</sup> | Percent of Large Carcasses Remaining After 20 Days | Mean Number of Days Large Carcasses Remained <sup>2,3</sup> | Percent of Small Carcasses Remaining After 20 Days | Mean Number of Days Small Carcasses Remained <sup>2,3</sup> |
|---|----------------------|--|---|--|---|--|---|
| August                                    | Grassland            | 33   | 21.0  | 0 <sup>4</sup>                                     | 7.0 <sup>4</sup>  | 50   | 35.0  |
| August                                    | CRP                  | 43   | 28.0  | 50   | 34.5  | 33   | 21.5  |
| Both Habitats                             |                      | 40   | 25.7  | 40   | 25.3  | 40   | 26.0  |
| September                                 | Grassland            | 0  | 2.3   | 0  | 3.0   | 0 <sup>4</sup>                                     | 1.0 <sup>4</sup>  |
| September                                 | CRP                  | 14   | 13.6  | 33   | 17.0  | 0  | 12.0  |
| Both Habitats                             |                      | 10   | 9.9   | 20   | 10.0  | 0  | 9.8   |
| October                                   | Grassland            | 33   | 21.0  | 50   | 35.0  | 0 <sup>4</sup>                                     | 7.0 <sup>4</sup>  |
| October                                   | CRP                  | 0  | 9.3   | 0  | 9.3   | 0  | 9.3   |
| Both Habitats                             |                      | 10   | 11.9  | 20   | 15.6  | 0  | 8.8   |
| Both Habitats Summer/Fall                 |                      | 20   | 14.6  | 27   | 16.3  | 13   | 13.2  |
| February                                  | Grassland            | 33   | 19.0  | 50   | 35.0  | 0 <sup>4</sup>                                     | 3.0 <sup>4</sup>  |
| February                                  | CRP                  | 0  | 6.6   | 0  | 9.3   | 0  | 4.5   |
| Both Habitats                             |                      | 10   | 9.3   | 20   | 15.6  | 0  | 4.2   |
| March                                     | Grassland            | 33   | 14.5  | 50   | 28.0  | 0 <sup>4</sup>                                     | 1.0 <sup>4</sup>  |
| March                                     | CRP                  | 14   | 10.3  | 33   | 22.5  | 0  | 4.3   |
| Both Habitats                             |                      | 20   | 11.4  | 40   | 24.3  | 0  | 3.6   |
| May                                       | Grassland            | 25   | 14.3  | 0  | 10.5  | 50   | 22.0  |
| May                                       | CRP                  | 0  | 5.3   | 0  | 6.3   | 0  | 4.3   |
| Both Habitats                             |                      | 10   | 8.3   | 0  | 8.0   | 20   | 8.8   |
| Both Habitats Winter/Spring               |                      | 13   | 9.62  | 20   | 14.7  | 7  | 5.29  |
| Overall Scavenger Removal Means and Rates |                      | 17   | 12.0  | 23   | 15.4  | 10   | 9.1   |

<sup>1</sup> CRP = Conservation Reserve Program land.

<sup>2</sup> Five ring-necked pheasants (representing large birds) and five cottontail quail (representing small birds and bats) were included in each scavenger removal trial.

<sup>3</sup> To calculate means, carcasses still present at the end of the 20-day carcass removal trial period were considered to have remained for 21 days.

<sup>4</sup> Sample size = one.

Table 3.7 Scavenger Removal Rates in Grassland Habitat and Conservation Reserve Program Land at Judith Gap Wind Energy Project.

| Habitat                       | Searcher Efficiency Trial Date | Percent of Carcasses Remaining After 20 Days | Mean Number of Days Carcasses Remained <sup>2,3</sup> | Percent of Large Carcasses Remaining After 20 Days | Mean Number of Days Large Carcasses Remained <sup>2,3</sup> | Percent of Small Carcasses Remaining After 20 Days | Mean Number of Days Small Carcasses Remained <sup>2,3</sup> |                  |
|-------------------------------|--------------------------------|--|---|--|---|--|---|------------------|
| <b>CRP<sup>1</sup></b>        | August                         | 43   | 28.0  | 50   | 34.5  | 33   | 21.5  |                  |
|                               | September                      | 14   | 13.6  | 33   | 17.0  | 0  | 12.0  |                  |
|                               | October                        | 0  | 9.3   | 0  | 9.3   | 0  | 9.3   |                  |
|                               | Subtotal Summer/Fall           | 19   | 15.2  | 30   | 18.7  | 9  | 12.8  |                  |
|                               | February                       | 0  | 6.6   | 0  | 9.3   | 0  | 4.5   |                  |
|                               | March                          | 14   | 10.3  | 33   | 22.5  | 0  | 4.3   |                  |
|                               | May                            | 0  | 5.3   | 0  | 6.3   | 0  | 4.3   |                  |
|                               | Subtotal Winter/Spring         | 5  | 7.4   | 11   | 11.5  | 0  | 4.4   |                  |
|                               | Total CRP                      | 12   | 11.1  | 21   | 14.9  | 5  | 8.4   |                  |
|                               | <b>Grassland</b>               | August                                       | 33  | 21.0   | 0.00 <sup>4</sup>   | 7.0 <sup>4</sup>                                   | 50  | 35.0             |
|                               |                                | September                                    | 0   | 2.3  | 0.00  | 3.0  | 0 <sup>4</sup>  | 1.0 <sup>4</sup> |
|                               |                                | October                                      | 33  | 21.0   | 50.00   | 35.0   | 0 <sup>4</sup>  | 7.0 <sup>4</sup> |
| Subtotal Summer/Fall          |                                | 22   | 13.0  | 20.00  | 12.0  | 25   | 14.3  |                  |
| February                      |                                | 33   | 19.0  | 50.00  | 35.0  | 0 <sup>4</sup>                                     | 3.0 <sup>4</sup>  |                  |
| March                         |                                | 33   | 14.5  | 50.00  | 28.0  | 0 <sup>4</sup>                                     | 1.0 <sup>4</sup>  |                  |
| May                           |                                | 25   | 14.3  | 0.00   | 10.5  | 50   | 22.0  |                  |
| Subtotal Winter/Spring        |                                | 30   | 15.7  | 33.33  | 21.0  | 250  | 8.7   |                  |
| Total Grassland               |                                | 26   | 14.4  | 27.27  | 16.5  | 25   | 11.5  |                  |
| <b>Both Habitats Combined</b> |                                |  | 17  | 12.0   | 23.33   | 15.4   | 10  | 9.1              |

<sup>1</sup> CRP = Conservation Reserve Program land.

<sup>2</sup> Five ring-necked pheasants (representing large birds) and five cottontail quail (representing small birds and bats) were included in each scavenger removal trial.

<sup>3</sup> To calculate means, carcasses still present at the end of the 20-day carcass removal trial period were considered to have remained for 21 days.

<sup>4</sup> Sample size = one.

---

Average carcass residency was shorter in CRP (14.9 days for large bird carcasses and 8.4 days for small bird carcasses) than in grassland (16.5 days for large bird carcasses and 11.5 days for small bird carcasses) (see Table 3.7).

### 3.5 TOTAL ESTIMATED FATALITIES FOR THE STUDY PERIOD

An average of 2.5 carcasses (15 birds and 35 bats) per turbine was found during carcass surveys at the 20 searched turbines during the study period. Searcher efficiency and scavenger removal rates were incorporated into estimates of bird and bat mortality. Because the low searcher efficiency for small birds (17%) was due at least in part to searcher efficiency trial carcasses being placed an average of 8.3 days before plot searches, and because small bird carcasses were estimated to remain on plots for an average of 9.1 days, an adjustment was made to the small bird searcher efficiency rate using the formula  $e^{PI/\bar{t}}$ , where PI = the mean interval between carcass placement and search date, and  $\bar{t}$  = mean carcass residency. This calculation accounted for the delay between carcass placement and search dates. Using the average small bird carcass residency (9.1 days) and adjusted searcher efficiency (42%) estimates for small birds, bat fatalities were estimated to be 13.4 bats/turbine (90% confidence interval = 8.04-24.72 bats/turbine) during the study period, or approximately 1,206 (90% confidence interval = 724-2,225) turbine-related bat fatalities during the study period at the 90 turbines (Table 3.8).

An average of 0.5 small birds (passerines) per turbine was found during searches. Using the 42% adjusted searcher efficiency rate and 9.1 mean carcass residency for small birds, small bird fatality rates were estimated to be 3.83 birds/turbine (90% confidence interval = 1.78-7.61), or a total of 344 small birds (90% confidence interval = 160-685) at the 90 turbines during the study period (see Table 3.8).

An average of 0.25 large birds (waterfowl and raptors) per turbine was found during carcass searches. Because the mean interval between searcher efficiency trial carcass placement and search dates was 7.2 days for large birds, and large bird carcasses remained an average of

Table 3.8 Estimated Avian and Bat Fatality Rates for JGWEP.<sup>1</sup>

| Category                 | Number of Carcasses Found | Estimated Total Mortality (90% Confidence Interval) | Standard Error | Carcasses/Turbine Estimate (90% Confidence Interval) | Standard Error |
|--------------------------|---------------------------|---|----------------|--|----------------|
| Small birds              | 10                        | 344 (160-685)                                       | 175            | 3.83 (1.78-7.61)                                     | 1.94           |
| Large birds <sup>2</sup> | 5                         | 62 (21-123)   | 32             | 0.69 (0.23-1.37)                                     | 0.36           |
| Raptors                  | 1                         | 12 (0-38)   | 13             | 0.14 (0-0.42)  | 0.14           |
| All Birds                | 15                        | 406 (28-593)  | 178            | 4.52 (0.31-6.59)                                     | 1.98           |
| Bats                     | 35                        | 1,206 (724-2,225)                                   | 491            | 13.40 (8.04-24.72)                                   | 5.46           |

<sup>1</sup> Project-wide estimates are rounded off to the nearest whole number.

<sup>2</sup> Includes one raptor.

15.43 days, it was assumed that the 77% searcher efficiency estimate for large birds was not inordinately impacted by the carcass placement-search date interval, and the large bird searcher efficiency rate was not adjusted. Using the carcass residency (15.43 days) and searcher efficiency (77%) for large birds, large bird fatalities were estimated at 0.69 large birds/turbine (90% confidence interval = 0.23-1.37), or a total of 62 large birds (90% confidence interval = 21-123) during the study period at the 90 turbines (see Table 3.8).

One raptor fatality was recorded during the study period, and using the large bird mean carcass residency (15.43 days) and searcher efficiency (77%) for large birds produces an estimate of 0.14 raptor fatalities/turbine (90% confidence interval = 0-0.42), or 14 raptors (90% confidence interval = 0-38) at the 90 turbines during the study period (see Table 3.8).

Estimated fatalities for small and large birds combined is 4.52 birds/turbine (90% confidence interval = 0.31-6.59), or 406 birds (90% confidence interval = 28-593) at the 90 turbines during the study period (see Table 3.8).

---

#### 4.0 DISCUSSION

Potential sources of bias are inevitably introduced into fatality studies. The species involved in turbine-related fatalities may not actually be scavenged at the same rate as the carcasses of species placed for scavenger removal trials. For example, the quail used during carcass trials may experience a higher or lower scavenging rate than carcasses normally occurring in the project area. Some studies found similar scavenger removal rates for bats and small birds (Young et al. 2003; Erickson et al. 2004; Fiedler 2004), but Johnson et al. (2003) found that bat carcasses remained an average of 10.36 days compared to 4.69 days for small bird carcasses. It is not known whether the small bird scavenger rates used to estimate carcass removal rates for bats accurately reflects the actual average carcass residency for bats at JGWEP. The inclusion in carcass totals of all carcasses found, including feather spots, may overestimate turbine-related mortality since some remains found may be due to natural predation; however, this may be offset to some extent by some turbine-related carcasses whose final location was outside the search area, either because they landed farther from the turbine, were moved by a scavenger, or the animal moved outside of search plots after being fatally injured. It seems likely that some turbine-related fatalities fell outside the search area because although searchers at JGWEP were 312 to 440 ft (95 to 134 m) from the tower when walking the search area boundaries, and 78% of the carcasses found were  $\leq 223$  ft (68 m) from the turbine, 16% were found  $\geq 318$  ft (97 m) from the search tower, with a range of 318 to 489 ft (97 to 149 m). It was also assumed that the 20 towers chosen for searching were representative of the project as a whole and do not collectively experience a fatality rate that leads to substantially inaccurate high or low fatality estimates for the entire turbine complex.

The mortality rates estimated for JGWEP are similar to those at other windfarms (Table 4.1), although the bat fatality rate was high compared to studies in the western U.S. Two windfarm studies from the eastern U.S. have reported higher bat fatality rates, but these wind farms were located on mountain ridgetops where migrating bats presumably were concentrated. Bat carcasses were found closer to turbines (average distance of 131 ft [40 m]) than bird carcasses (average distance of 210 ft [64 m]). This pattern has been noted in other studies (Erickson et al. 2000; Erickson et al. 2003; Erickson et al. 2004; Jain et al. 2007).

Table 4.1 Estimated Fatality Rates for Birds and Bats at Other U.S. Wind Farms and JGWEP.<sup>1</sup>

| Location                          | Bird Fatalities/<br>Turbine/Year | Bat Fatalities/<br>Turbine/Year | Searcher Efficiency Birds<br>(%) (Large, Medium,<br>Small) | Carcass Residency<br>(Days) (Large, Medium,<br>Small) |
|-----------------------------------|----------------------------------|---------------------------------|--|---|
| Vansycle, OR <sup>2</sup>         | 0.63                             | 0.74                            | 88 L, 50 S   | 25  |
| Nine Canyon, WA <sup>3</sup>      | 3.59                             | 3.21                            | 78 L, 44 S   | 33 L, 11 S  |
| Klondike, OR <sup>4</sup>         | 1.42                             | 1.16                            | 92 L, 75 S   | 19.9 L, 14.2 S  |
| Buffalo Ridge, MN <sup>5</sup>    | 0.98-4.45                        | n/a                             | 49 L, 40 M, 29 S   | 8.5 L, 8 M, 4.7 S                                     |
| Mountaineer, WV <sup>6</sup>      | 4.04                             | 47.53                           | 28 (Combined)  | 6.7 (Combined)  |
| Stateline, OR <sup>7</sup>        | 1.93                             | 1.12                            | 78 L, 42 S   | 35.7 L/M, 16.7 S                                      |
| Foot Creek Rim, WY <sup>8</sup>   | 1.5                              | 1.34                            | 92L, 87 M, 59 S  | 29.5 L, 37.3 M, 13.4 S                                |
| Buffalo Mountain, TN <sup>9</sup> | 1.8                              | 63.9                            | Variable   | Variable  |
| Judith Gap, MT <sup>10</sup>      | 4.52                             | 13.4                            | 77 L, 17 S   | 15.4 L, 9.1 S   |

<sup>1</sup> All fatality rates have incorporated searcher efficiency and scavenger removal data. Studies of <1 year duration are indicated.

<sup>2</sup> Erickson et al. (2000).

<sup>3</sup> Erickson et al. (2003).

<sup>4</sup> Johnson et al. (2003).

<sup>5</sup> Johnson et al. (2000).

<sup>6</sup> Kerns and Kerlinger (2004); 7-month study.

<sup>7</sup> Erickson et al. (2004).

<sup>8</sup> Young et al. (2003).

<sup>9</sup> Fiedler et al. (2007); 9-month study.

<sup>10</sup> This study; 7-month study.

Ninety-seven percent of all bat carcasses were found during fall migration. Johnson (2004) noted that wind farms pose the greatest risk to bats during fall migration, and the timing of bat mortality events at JGWEP, as well as species composition (highly migratory, tree-roosting species) (Cryan and Brown 2007), is similar to that in other studies (Erickson et al. 2000; Erickson et al. 2003; Erickson et al. 2004; Fiedler et al. 2007; Kerns and Kerlinger 2004; Jain et al. 2007).

Because of the dates on which they were found, waterfowl fatalities likely represented migrating birds. Although several small reservoirs and small waterways associated with Roberts Creek and its tributaries occur within 0.6 mi (1.0 km) of the project area, these are used only by mallards and other dabbling ducks and by occasional migrating shorebirds (personal communication,

August 13, 2007, with Ralph Rogers, Centmont Bioconsultants). Waterfowl (including American coots) made up 50% of the total bird fatalities recorded during the study (see Table 3.1), but waterfowl are considerably larger than passerines and their carcasses are more likely to be discovered incidentally. By contrast, horned larks (39%) and McCown's longspurs (41%) together comprised 80% of the birds detected during grassland bird surveys in 2003 and 2007 (Appendix A, Table A.3) but only 27% (all horned larks) of the carcasses found during turbine plot searches (see Table 3.2).

The results of this study suggest that avian fatality rates at JGWEP are similar to fatality rates at other wind plants around the U.S. (see Table 4.1--note that some studies were conducted for <1 year), whereas estimated fatality rates for bats are somewhat higher when compared to other studies in the western U.S.

---

## 5.0 REFERENCES

- Cryan, P.M., and A.C Brown. 2007. Migration of bats past a remote island offers clues toward the problem of bat fatalities at wind turbines. *Biological Conservation* 139:1-11.
- Erickson, W.P., G.D. Johnson, M.D. Strickland, and K. Kronner. 2000. Avian and bat mortality associated with the Vansycle Wind Project, Umatilla County, Oregon: 1999 study year. Technical Report prepared for Umatilla County Department of Resource Services and Development, Pendleton, Oregon.
- Erickson, W.P., B. Gritski, and K. Kronner. 2003. Nine Canyon Wind Power Project avian and bat monitoring report, September 2002-August 2003. Technical report submitted to Energy Northwest and the Nine Canyon Technical Advisory Committee.
- Erickson, W.P., J. Jeffrey, K. Kronner, and K. Bay. 2004. Stateline Wind Project Wildlife Monitoring Final Report, July 2001-December 2003. Technical report submitted to FPL Energy, the Oregon Energy Facility Siting Council, and the Stateline Technical Advisory Committee.
- Fiedler, J.K. 2004. Assessment of bat mortality and activity at Buffalo Mountain Windfarm, eastern Tennessee. Master's Thesis, University of Tennessee, Knoxville.
- Fiedler, J.K., T.H. Henry, R.D. Tankersley, and C.P. Nicholson. 2007. Results of bat and bird mortality monitoring at the expanded Buffalo Mountain windfarm, 2005. Tennessee Valley Authority, Knoxville.
- Higgins, K.F., R.G. Osborn, C.D. Dieter, and R.E. Usgaard. 1996. Monitoring of seasonal bird activity and mortality at the Buffalo Ridge Wind Resource Area, Minnesota, 1994-1995. Completion report for the research period May 1, 1994-December 31, 1995. Unpublished report prepared for Kenetech Windpower, Inc. by the South Dakota Cooperative Fish and Wildlife Research Unit, Brookings, South Dakota. 84 pp.
- Howell, J.A., and J.E. DiDonato. 1991. Assessment of avian use and mortality related to wind turbine operations, Altamont Pass, Alameda and Contra Costa Counties, California. September 1988 through August 1989. Final report. Prepared for U.S. Windpower, Inc., Livermore, California. 72 pp.
- Jain, A., P. Kerlinger, R. Curry, and L. Slobodnik. 2007. Annual report for the Maple Ridge Wind Power Project post-construction bird and bat fatality study-2006. Prepared for PPM Energy and Horizon Energy and Technical Advisory Committee for the Maple Ridge Project Study by Curry and Kerlinger, LLC. 53 pp.

- 
- Johnson, G. 2004. A review of bat impacts at wind farms in the U.S. In Proceedings of the wind energy birds/bats workshop: Understanding and resolving bird and bat impacts, Washington, DC, May 18-19, 2004 (S.S. Schwartz, ed.). Prepared by RESOLVE, Inc., Washington, D.C. 107 pp. + append.
- Johnson, G.D., W.P. Erickson, M.D. Strickland, M.F. Shepherd, and D.S. Shepherd. 2000. Avian monitoring studies at the Buffalo Ridge, Minnesota Wind Resource Area: results of a 4-year study. Western Ecosystems Technology, Inc., Cheyenne, Wyoming.
- Johnson, G.D., W.P. Erickson, M.D. Strickland, M.F. Shepherd, D.A. Shepherd, and S.A. Sarappo. 2002. Collision mortality of local and migrant birds at a large-scale wind power development on Buffalo Ridge, Minnesota. *Wildlife Society Bulletin* 30:879-887.
- Johnson, G., W. Erickson, J. White, and R. McKinney. 2003. Avian and bat mortality during the first year of operation at the Klondike Phase I Wind Project, Sherman County, Oregon. Prepared for Northwestern Wind Power by WEST, Inc., Cheyenne, Wyoming. 17 pp.
- Kerns, J., and P. Kerlinger. 2004. A study of bird and bat collision fatalities at the Mountaineer Wind Energy Center, Tucker County, West Virginia: annual report for 2003. Prepared for FPL Energy and Mountaineer Wind Energy Center Technical Review Committee by Curry and Kerlinger, LLC.
- Leddy, K.L., K.F. Higgins, and D.E. Naugle. 1999. Effects of wind turbines on upland nesting birds in conservation reserve program grasslands. *Wilson Bulletin* 111(1):100-104.
- McCrary, M.D., R.L. McKernan, and R.W. Schreiber. 1986. San Geronio wind resource area: Impacts of commercial wind turbine generators on birds, 1985 data report. Prepared for Southern California Edison Company. 33 pp.
- Morrison, M. 2002. Searcher bias and scavenging rates in bird/wind energy studies. National Renewable Energy Laboratory, Golden, Colorado. NREL/SR-500-30876. 5 pp.
- Orloff, S., and A. Flannery. 1992. Wind turbine effects on avian activity, habitat use, and mortality in Altamont Pass and Solano County Wind Resource Areas, 1989-1991. Final report. Prepared for the Planning Departments of Alameda, Contra Costa, and Solano Counties and the California Energy Commission.
- Strickland, D. 2004. Non-fatality and habitat impacts on birds from wind energy development. *In* Proceedings of the wind energy birds/bats workshop: Understanding and resolving bird and bat impacts, Washington, D.C., May 18-19, 2004 (S.S. Schwartz, ed.). Prepared by RESOLVE, Inc., Washington, D.C. 107 pp. + append.
- TRC Mariah Associates Inc. 2006. Avian and bat monitoring plan, Judith Gap Wind Energy Project, Wheatland County, Montana. Prepared for Judith Gap Energy, LLC, by TRC Mariah Associates Inc. 17 pp.

- 
- U.S. Farm Service Agency and Montana Department of State Lands. 2004. Environmental assessment for the Judith Gap wind farm project. Montana Department of Natural Resources and Conservation, Real Estate Management Bureau, Helena, Montana, and U.S. Farm Service Agency, Bozeman, Montana. 51 pp. + append.
- Young, D.P., Jr., W.P. Erickson, R.E. Good, M.D. Strickland, and G.D. Johnson. 2003. Avian and bat mortality associated with the initial phase of the Foote Creek Rim Windpower Project, Carbon County, Wyoming, November 1998-June 2002. Final Report. Prepared for PacifiCorp, Inc., Portland, Oregon; SeaWest Windpower Inc., San Diego, California; and Bureau of Land Management, Rawlins District Office, Rawlins, Wyoming, by Western Ecosystems Technology, Inc., Cheyenne, Wyoming.

**APPENDIX A:**  
GRASSLAND BIRD DISPLACEMENT STUDY

## **1.0 INTRODUCTION**

Wind energy plants constructed in grassland habitats have the potential to displace grassland breeding birds in the vicinity of the turbines (Leddy et al. 1999; Johnson et al. 2000; Erickson et al. 2004). To examine the impact of the Judith Gap Wind Energy Project (JGWEP) on breeding grassland birds, point count surveys that were conducted in 2003 prior to construction activities (Flath 2003) were repeated in 2007 after the wind farm was completed and became operational. Comparisons were made between numbers of birds recorded during the pre- and post-construction bird surveys to determine if the construction of the wind plant had any detectable impact on locally breeding grassland birds.

---

## 2.0 METHODS

Flath (2003) established 13 point count transects in each of the habitat types found on the wind project: natural grasslands, wheat stubble, and Conservation Reserve Program (CRP) (Figure A.1). Eight of the 13 transects were within the future footprint of the windfarm area, and five control transects were established outside the proposed windfarm in habitat types similar to those found within the windfarm. The number of transects in each habitat type approximated the relative percentage of the various habitat types in the study area. Each 0.78-mi (1.25-km) transect consisted of six point count locations separated by 820 ft (250 m). Flath (2003) counted each bird seen or heard within a 328-ft (100-m) radius circle from each point during a 10-minute period. Counts were conducted between 6:45 a.m. (after the dawn chorus ended) and 10:00 a.m. Counts on all transects were conducted during the breeding season in the last two weeks of April, May, and June. A “survey” was defined as the completion of one round of point counts at all six points on each of the 13 transects.

Flath’s (2003) study was repeated, using the same transects and methodology, during the spring of 2007 after all wind turbines in the project area were fully operational. The beginning and end points of each transect were recorded using a global positioning system (GPS) unit, and distances between survey points were traversed on an all-terrain vehicle. In 2007, transects 1-8 (the experimental transects) were immediately adjacent to recently constructed wind towers, and transects 9-13 (the control transects) were outside the wind plant ( $\geq 1.0$  mi [1.6 km] from the nearest turbine) in habitats unchanged since the 2003 surveys.

Only grassland birds with  $\geq 5$  total detections during the six surveys (long-billed curlew; mountain plover; horned lark; Sprague’s pipit; grasshopper, savannah, and vesper sparrows; McCown’s longspur; western meadowlark) were used for analysis. Potential differences in bird numbers between the 2003 and 2007 experimental and control plots were examined using two-sample t-tests. Differences in birds/transect between years were calculated for each species for both experimental and control transects by subtracting 2003 numbers from 2007 numbers for each transect, after which two-sample t-tests were performed on the differences to detect any significant changes in numbers between years ( $\alpha = 0.05$ ).




Figure A.1 Wind Turbine Arrays and Grassland Bird Displacement Study Transects.

### **3.0 RESULTS**

Bird numbers recorded during point count surveys increased between 2003 and 2007 in both the experimental and control areas for six of the nine grassland bird species analyzed (Table A.1). Overall totals for grassland birds were 54% higher on the experimental transects and 20% higher on the control transects in 2007 compared to 2003. The results of the t-tests indicated significant increases for two species (western meadowlark,  $p = 0.048$ ; vesper sparrow,  $p = 0.044$ ), but a comparison of overall bird numbers indicated no significant change ( $p = 0.072$ ). Table A.2 presents the combined counts for both years for all birds, and Table A.3 presents the results of each of the six surveys.

Table A.1 Results of Monthly (April, May, and June) Point Counts for Grassland Birds on Experimental and Control Transects at Judith Gap Wind Energy Project.

| Species                                      | 2003 | 2007  | Difference (% Change) |
|--|------|-------|-----------------------|
| <b>Experimental Transects (n = 8/survey)</b> |      |       |                       |
| Mountain plover                              | 2    | 18    | 16 (900)              |
| Long-billed curlew                           | 0    | 2     | 2 (n/a)               |
| Horned lark                                  | 393  | 502   | 109 (28)              |
| Sprague's pipit                              | 29   | 21    | -8 (-28)              |
| Grasshopper sparrow                          | 5    | 0     | -5 (n/a)              |
| Vesper sparrow                               | 13   | 85    | 72 (554) <sup>1</sup> |
| Savannah sparrow                             | 75   | 77    | 2 (3)                 |
| McCown's longspur                            | 330  | 562   | 232 (70)              |
| Western meadowlark                           | 37   | 93    | 56 (151) <sup>1</sup> |
| Totals                                       | 884  | 1,360 | 476 (54)              |
| <b>Control Transects (n = 5/survey)</b>      |      |       |                       |
| Mountain plover                              | 0    | 0     | 0 (0)                 |
| Long-billed curlew                           | 13   | 17    | 4 (31)                |
| Horned lark                                  | 264  | 258   | -6 (-2)               |
| Sprague's pipit                              | 2    | 20    | 18 (900)              |
| Grasshopper sparrow                          | 0    | 0     | 0 (0)                 |
| Vesper sparrow                               | 2    | 5     | 3 (150)               |
| Savannah sparrow                             | 43   | 48    | 5 (12)                |
| McCown's longspur                            | 257  | 344   | 87 (34)               |
| Western meadowlark                           | 10   | 17    | 7 (70)                |
| Totals                                       | 591  | 709   | 118 (20)              |

<sup>1</sup> Significant at  $\alpha = 0.05$  level.

Table A.2 Total Numbers of Various Species Detected During Point Counts Conducted at JGWEP in 2003 and 2007.

| Species              | 2003  | 2007  | Total | % of Total      |
|----------------------|-------|-------|-------|-----------------|
| McCown's longspur    | 587   | 906   | 1,493 | 41              |
| Horned lark          | 657   | 760   | 1,417 | 39              |
| Savannah sparrow     | 118   | 125   | 243   | 7               |
| Western meadowlark   | 47    | 110   | 157   | 4               |
| Vesper sparrow       | 15    | 90    | 105   | 3               |
| Sprague's pipit      | 31    | 41    | 72    | 2               |
| Long-billed curlew   | 13    | 19    | 32    | 1               |
| Mountain plover      | 2     | 18    | 20    | 1               |
| Gadwall              | 0     | 13    | 13    | 0               |
| Northern pintail     | 2     | 8     | 10    | 0               |
| Mallard              | 0     | 8     | 8     | 0               |
| Northern harrier     | 3     | 5     | 8     | 0               |
| Grasshopper sparrow  | 5     | 0     | 5     | 0               |
| Red-winged blackbird | 0     | 5     | 5     | 0               |
| Gray partridge       | 3     | 2     | 5     | 0               |
| Sandhill crane       | 0     | 5     | 5     | 0               |
| Ferruginous hawk     | 0     | 4     | 4     | 0               |
| Killdeer             | 0     | 4     | 4     | 0               |
| Brown-headed cowbird | 0     | 3     | 3     | 0               |
| Barn swallow         | 0     | 2     | 2     | 0               |
| Chipping sparrow     | 2     | 0     | 2     | 0               |
| Lark bunting         | 2     | 0     | 2     | 0               |
| American kestrel     | 1     | 0     | 1     | 0               |
| Northern flicker     | 0     | 1     | 1     | 0               |
| American robin       | 0     | 1     | 1     | 0               |
| Ring-billed gull     | 0     | 1     | 1     | 0               |
| Clay-colored sparrow | 0     | 1     | 1     | 0               |
| Canada goose         | 0     | 1     | 1     | 0               |
| American coot        | 0     | 1     | 1     | 0               |
| Total                | 1,488 | 2,134 | 3,622 | 98 <sup>1</sup> |

<sup>1</sup> Total does not equal 100 due to rounding errors.

Table A.3 Results of Point Counts Conducted at JGWEC in 2003 and 2007.<sup>1</sup>

| Species              | T1 | T2 | T3 | T4 | T5 | T6 | T7 | T8 | Experimental | T9 | T10 | T11 | T12 | T13 | Control | Total |
|----------------------|----|----|----|----|----|----|----|----|--------------|----|-----|-----|-----|-----|---------|-------|
| <b>April 2003</b>    |    |    |    |    |    |    |    |    |              |    |     |     |     |     |         |       |
| Horned lark          | 12 | 24 | 20 | 35 | 28 | 7  | 8  | 4  | 138          | 6  | 25  | 1   | 21  | 24  | 77      | 215   |
| McCown's longspur    | 48 | 9  | 25 | 0  | 41 | 0  | 1  | 5  | 129          | 43 | 38  | 0   | 2   | 20  | 103     | 232   |
| Western meadowlark   | 0  | 0  | 0  | 0  | 0  | 9  | 8  | 2  | 19           | 0  | 0   | 5   | 0   | 0   | 5       | 24    |
| Vesper sparrow       | 0  | 0  | 0  | 0  | 0  | 0  | 1  | 0  | 1            | 0  | 0   | 0   | 0   | 0   | 0       | 1     |
| Savannah sparrow     | 0  | 0  | 0  | 0  | 0  | 2  | 0  | 0  | 2            | 0  | 0   | 2   | 0   | 0   | 2       | 4     |
| Brown-headed cowbird | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Mountain plover      | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Sprague's pipit      | 0  | 0  | 0  | 0  | 0  | 1  | 3  | 4  | 8            | 0  | 0   | 0   | 0   | 0   | 0       | 8     |
| Northern harrier     | 0  | 0  | 0  | 0  | 0  | 1  | 0  | 0  | 1            | 1  | 0   | 0   | 0   | 0   | 1       | 2     |
| Long-billed curlew   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 2   | 0   | 0   | 0   | 2       | 2     |
| Killdeer             | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Ferruginous hawk     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Red-winged blackbird | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Northern flicker     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Sandhill crane       | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Gray partridge       | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Mallard              | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| American robin       | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Ring-billed gull     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Clay-colored sparrow | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Northern pintail     | 0  | 0  | 0  | 0  | 0  | 2  | 0  | 0  | 2            | 0  | 0   | 0   | 0   | 0   | 0       | 2     |
| Gadwall              | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Canada goose         | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |

Table A.3 (Continued)

| Species              | T1 | T2 | T3 | T4 | T5 | T6 | T7 | T8 | Experimental | T9 | T10 | T11 | T12 | T13 | Control | Total |
|----------------------|----|----|----|----|----|----|----|----|--------------|----|-----|-----|-----|-----|---------|-------|
| American kestrel     | 0  | 0  | 0  | 1  | 0  | 0  | 0  | 0  | 1            | 0  | 0   | 0   | 0   | 0   | 0       | 1     |
| Grasshopper sparrow  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Lark bunting         | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Barn swallow         | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| American coot        | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| <b>Totals</b>        |    |    |    |    |    |    |    |    |              |    |     |     |     |     |         | 491   |
| <b>May 2003</b>      |    |    |    |    |    |    |    |    |              |    |     |     |     |     |         | 301   |
| Horned lark          | 22 | 16 | 21 | 21 | 17 | 7  | 10 | 0  | 114          | 7  | 13  | 0   | 32  | 24  | 76      | 190   |
| McCown's longspur    | 31 | 9  | 31 | 4  | 21 | 1  | 0  | 0  | 97           | 23 | 39  | 0   | 6   | 11  | 79      | 176   |
| Western meadowlark   | 0  | 1  | 1  | 0  | 0  | 3  | 4  | 2  | 11           | 1  | 0   | 0   | 0   | 1   | 2       | 13    |
| Vesper sparrow       | 0  | 0  | 0  | 0  | 0  | 2  | 3  | 5  | 10           | 0  | 0   | 0   | 0   | 0   | 0       | 10    |
| Savannah sparrow     | 0  | 0  | 0  | 2  | 0  | 15 | 3  | 17 | 37           | 0  | 1   | 17  | 0   | 1   | 19      | 56    |
| Brown-headed cowbird | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Mountain plover      | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Sprague's pipit      | 0  | 0  | 0  | 0  | 0  | 1  | 4  | 6  | 11           | 0  | 2   | 0   | 0   | 0   | 2       | 13    |
| Northern harrier     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Long-billed curlew   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 1  | 2   | 0   | 0   | 0   | 3       | 3     |
| Killdeer             | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Ferruginous hawk     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Red-winged blackbird | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Northern flicker     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Sandhill crane       | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Gray partridge       | 0  | 0  | 0  | 0  | 0  | 1  | 0  | 0  | 1            | 0  | 0   | 2   | 0   | 0   | 2       | 3     |
| Mallard              | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |

Table A.3 (Continued)

| Species              | T1 | T2 | T3 | T4 | T5 | T6 | T7 | T8 | Experimental | T9 | T10 | T11 | T12 | T13 | Control | Total |
|----------------------|----|----|----|----|----|----|----|----|--------------|----|-----|-----|-----|-----|---------|-------|
| American robin       | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Ring-billed gull     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Clay-colored sparrow | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Northern pintail     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Gadwall              | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Canada goose         | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| American kestrel     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Chipping sparrow     | 0  | 0  | 0  | 2  | 0  | 0  | 0  | 0  | 2            | 0  | 0   | 0   | 0   | 0   | 0       | 2     |
| Grasshopper sparrow  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Lark bunting         | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Barn swallow         | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| American coot        | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| <b>Totals</b>        |    |    |    |    |    |    |    |    |              |    |     |     |     |     |         | 466   |
| <b>June 2003</b>     |    |    |    |    |    |    |    |    |              |    |     |     |     |     |         | 283   |
| Horned lark          | 20 | 17 | 18 | 30 | 29 | 5  | 22 | 0  | 141          | 11 | 36  | 0   | 36  | 28  | 111     | 252   |
| McCown's longspur    | 34 | 11 | 42 | 1  | 16 | 0  | 0  | 0  | 104          | 36 | 21  | 0   | 13  | 5   | 75      | 179   |
| Western meadowlark   | 0  | 0  | 0  | 0  | 0  | 2  | 1  | 4  | 7            | 0  | 0   | 1   | 1   | 1   | 3       | 10    |
| Vesper sparrow       | 0  | 0  | 1  | 0  | 0  | 0  | 1  | 0  | 2            | 0  | 0   | 0   | 0   | 2   | 2       | 4     |
| Savannah sparrow     | 0  | 1  | 0  | 0  | 0  | 20 | 8  | 7  | 36           | 1  | 1   | 17  | 0   | 3   | 22      | 58    |
| Brown-headed cowbird | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |
| Mountain plover      | 0  | 0  | 0  | 0  | 2  | 0  | 0  | 0  | 2            | 0  | 0   | 0   | 0   | 0   | 0       | 2     |
| Sprague's pipit      | 0  | 0  | 0  | 0  | 0  | 1  | 4  | 5  | 10           | 0  | 0   | 0   | 0   | 0   | 0       | 10    |
| Northern harrier     | 0  | 0  | 0  | 1  | 0  | 0  | 0  | 0  | 1            | 0  | 0   | 0   | 0   | 0   | 0       | 1     |
| Long-billed curlew   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 8   | 0   | 0   | 0   | 8       | 8     |

Table A.3 (Continued)

| Species              | T1  | T2 | T3 | T4 | T5 | T6 | T7 | T8 | Experimental | T9 | T10 | T11 | T12 | T13 | Control | Total |     |
|----------------------|-----|----|----|----|----|----|----|----|--------------|----|-----|-----|-----|-----|---------|-------|-----|
| Killdeer             | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Ferruginous hawk     | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Red-winged blackbird | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Northern flicker     | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Sandhill crane       | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Gray partridge       | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Mallard              | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| American robin       | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Ring-billed gull     | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Clay-colored sparrow | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Northern pintail     | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Gadwall              | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Canada goose         | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| American kestrel     | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Chipping sparrow     | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Grasshopper sparrow  | 0   | 0  | 0  | 0  | 0  | 4  | 1  | 0  | 5            | 0  | 0   | 0   | 0   | 0   | 0       | 5     |     |
| Lark bunting         | 0   | 0  | 1  | 0  | 0  | 0  | 0  | 0  | 1            | 0  | 0   | 1   | 0   | 0   | 1       | 2     |     |
| Barn swallow         | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| American coot        | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 0   | 0   | 0   | 0       | 0     |     |
| Totals               | 309 |    |    |    |    |    |    |    |              |    |     |     |     |     |         | 222   | 531 |

**April 2007**

|                    |    |    |    |    |    |    |    |   |     |    |    |    |    |    |     |     |
|--------------------|----|----|----|----|----|----|----|---|-----|----|----|----|----|----|-----|-----|
| Horned lark        | 12 | 39 | 10 | 58 | 26 | 13 | 20 | 9 | 187 | 19 | 22 | 13 | 22 | 22 | 98  | 285 |
| McCown's longspur  | 42 | 4  | 25 | 2  | 40 | 0  | 0  | 9 | 122 | 49 | 51 | 0  | 5  | 20 | 125 | 247 |
| Western meadowlark | 0  | 2  | 4  | 2  | 2  | 6  | 7  | 5 | 28  | 0  | 1  | 5  | 0  | 0  | 6   | 34  |
| Vesper sparrow     | 0  | 0  | 0  | 0  | 0  | 0  | 2  | 0 | 2   | 0  | 0  | 0  | 0  | 0  | 0   | 2   |

Table A.3 (Continued)

| Species              | T1 | T2 | T3 | T4 | T5 | T6 | T7 | T8 | Experimental | T9 | T10 | T11 | T12 | T13 | Control | Total |
|----------------------|----|----|----|----|----|----|----|----|--------------|----|-----|-----|-----|-----|---------|-------|
| Savannah sparrow     | 0  | 0  | 0  | 0  | 0  | 1  | 1  | 0  | 2            | 0  | 0   | 4   | 0   | 0   | 4       | 6     |
| Mountain plover      | 2  | 0  | 0  | 0  | 4  | 0  | 0  | 0  | 6            | 0  | 0   | 0   | 0   | 0   | 0       | 6     |
| Sprague's pipit      | 0  | 0  | 0  | 0  | 0  | 4  | 3  | 0  | 7            | 0  | 3   | 0   | 0   | 0   | 3       | 10    |
| Northern harrier     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 1  | 1            | 1  | 0   | 0   | 0   | 0   | 1       | 2     |
| Long-billed curlew   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 4  | 2   | 0   | 0   | 0   | 6       | 6     |
| Ferruginous hawk     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 1   | 0   | 0   | 1       | 1     |
| Red-winged blackbird | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 2   | 0   | 0   | 2       | 2     |
| Northern flicker     | 0  | 0  | 1  | 0  | 0  | 0  | 0  | 0  | 1            | 0  | 0   | 0   | 0   | 0   | 0       | 1     |
| Sandhill crane       | 0  | 0  | 5  | 0  | 0  | 0  | 0  | 0  | 5            | 0  | 0   | 0   | 0   | 0   | 0       | 5     |
| Gray partridge       | 0  | 0  | 0  | 2  | 0  | 0  | 0  | 0  | 2            | 0  | 0   | 0   | 0   | 0   | 0       | 2     |
| Mallard              | 0  | 0  | 0  | 2  | 0  | 0  | 0  | 0  | 2            | 6  | 0   | 0   | 0   | 0   | 6       | 8     |
| American robin       | 0  | 0  | 0  | 0  | 1  | 0  | 0  | 0  | 1            | 0  | 0   | 0   | 0   | 0   | 0       | 1     |
| Ring-billed gull     | 0  | 0  | 0  | 0  | 1  | 0  | 0  | 0  | 1            | 0  | 0   | 0   | 0   | 0   | 0       | 1     |
| Clay-colored sparrow | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 1  | 1            | 0  | 0   | 0   | 0   | 0   | 0       | 1     |
| Northern pintail     | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 8  | 0   | 0   | 0   | 0   | 8       | 8     |
| Gadwall              | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 10 | 0   | 0   | 0   | 0   | 10      | 10    |
| Canada goose         | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 1   | 0   | 0   | 0   | 1       | 1     |
| Totals               | 56 | 45 | 45 | 66 | 74 | 24 | 33 | 25 | 368          | 97 | 80  | 25  | 27  | 42  | 271     | 639   |
| <b>May 2007</b>      |    |    |    |    |    |    |    |    |              |    |     |     |     |     |         |       |
| Horned lark          | 18 | 26 | 23 | 31 | 24 | 12 | 28 | 0  | 162          | 7  | 20  | 0   | 28  | 24  | 79      | 241   |
| McCown's longspur    | 67 | 16 | 60 | 27 | 23 | 0  | 0  | 15 | 208          | 21 | 33  | 0   | 2   | 22  | 78      | 286   |
| Western meadowlark   | 2  | 0  | 5  | 2  | 4  | 3  | 5  | 13 | 34           | 1  | 3   | 0   | 0   | 1   | 5       | 39    |
| Vesper sparrow       | 0  | 0  | 16 | 0  | 2  | 10 | 11 | 5  | 44           | 0  | 0   | 0   | 0   | 2   | 2       | 46    |

Table A.3 (Continued)

| Species              | T1 | T2 | T3  | T4 | T5 | T6 | T7 | T8 | Experimental | T9 | T10 | T11 | T12 | T13 | Control | Total |
|----------------------|----|----|-----|----|----|----|----|----|--------------|----|-----|-----|-----|-----|---------|-------|
| Savannah sparrow     | 0  | 0  | 1   | 0  | 0  | 11 | 2  | 17 | 31           | 0  | 0   | 21  | 0   | 1   | 22      | 53    |
| Brown-headed cowbird | 0  | 0  | 3   | 0  | 0  | 0  | 0  | 0  | 3            | 0  | 0   | 0   | 0   | 0   | 0       | 3     |
| Mountain plover      | 0  | 0  | 0   | 2  | 4  | 2  | 0  | 0  | 8            | 0  | 0   | 0   | 0   | 0   | 0       | 8     |
| Sprague's pipit      | 0  | 0  | 0   | 0  | 0  | 0  | 0  | 8  | 8            | 0  | 8   | 0   | 0   | 0   | 8       | 16    |
| Northern harrier     | 0  | 0  | 0   | 0  | 0  | 0  | 1  | 0  | 1            | 0  | 0   | 0   | 0   | 0   | 0       | 1     |
| Long-billed curlew   | 0  | 0  | 0   | 0  | 0  | 0  | 0  | 0  | 0            | 3  | 1   | 0   | 0   | 0   | 4       | 4     |
| Killdeer             | 0  | 0  | 0   | 0  | 0  | 0  | 0  | 0  | 0            | 2  | 0   | 0   | 0   | 0   | 2       | 2     |
| Ferruginous hawk     | 0  | 0  | 0   | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 1   | 0   | 0   | 0   | 1       | 1     |
| Red-winged blackbird | 0  | 0  | 0   | 0  | 0  | 0  | 0  | 0  | 0            | 0  | 0   | 2   | 0   | 0   | 2       | 2     |
| Totals               | 87 | 42 | 108 | 62 | 57 | 38 | 47 | 58 | 499          | 34 | 66  | 23  | 30  | 50  | 203     | 702   |
| <b>June 2007</b>     |    |    |     |    |    |    |    |    |              |    |     |     |     |     |         |       |
| Horned lark          | 25 | 33 | 21  | 14 | 20 | 10 | 20 | 10 | 153          | 2  | 17  | 0   | 38  | 24  | 81      | 234   |
| McCown's longspur    | 48 | 12 | 58  | 40 | 72 | 0  | 0  | 2  | 232          | 66 | 58  | 0   | 10  | 7   | 141     | 373   |
| Western meadowlark   | 2  | 0  | 1   | 0  | 6  | 7  | 5  | 10 | 31           | 3  | 0   | 2   | 0   | 1   | 6       | 37    |
| Vesper sparrow       | 0  | 0  | 6   | 0  | 0  | 11 | 3  | 19 | 39           | 0  | 0   | 0   | 0   | 3   | 3       | 42    |
| Savannah sparrow     | 0  | 0  | 0   | 0  | 0  | 13 | 5  | 26 | 44           | 4  | 0   | 15  | 0   | 3   | 22      | 66    |
| Mountain plover      | 0  | 0  | 0   | 1  | 3  | 0  | 0  | 0  | 4            | 0  | 0   | 0   | 0   | 0   | 0       | 4     |
| Sprague's pipit      | 0  | 0  | 0   | 0  | 0  | 4  | 0  | 2  | 6            | 2  | 7   | 0   | 0   | 0   | 9       | 15    |
| Northern harrier     | 0  | 0  | 0   | 0  | 0  | 0  | 1  | 0  | 1            | 0  | 1   | 0   | 0   | 0   | 1       | 2     |
| Long-billed curlew   | 0  | 0  | 0   | 0  | 0  | 0  | 0  | 2  | 2            | 5  | 2   | 0   | 0   | 0   | 7       | 9     |
| Killdeer             | 0  | 0  | 0   | 0  | 0  | 0  | 0  | 0  | 0            | 2  | 0   | 0   | 0   | 0   | 2       | 2     |
| Ferruginous hawk     | 0  | 0  | 0   | 1  | 0  | 0  | 0  | 0  | 1            | 0  | 1   | 0   | 0   | 0   | 1       | 2     |

Table A.3 (Continued)

| Species              | T1 | T2 | T3 | T4 | T5  | T6 | T7 | T8 | Experimental | T9 | T10 | T11 | T12 | T13 | Control | Total |
|----------------------|----|----|----|----|-----|----|----|----|--------------|----|-----|-----|-----|-----|---------|-------|
| Red-winged blackbird | 0  | 0  | 0  | 0  | 0   | 0  | 0  | 0  | 0            | 1  | 0   | 0   | 0   | 0   | 1       | 1     |
| Gadwall              | 0  | 0  | 0  | 0  | 0   | 0  | 0  | 0  | 0            | 3  | 0   | 0   | 0   | 0   | 3       | 3     |
| Barn swallow         | 0  | 0  | 0  | 0  | 0   | 0  | 0  | 0  | 0            | 2  | 0   | 0   | 0   | 0   | 2       | 2     |
| American coot        | 0  | 0  | 0  | 0  | 0   | 0  | 0  | 0  | 0            | 1  | 0   | 0   | 0   | 0   | 1       | 1     |
| Totals               | 75 | 45 | 86 | 56 | 101 | 45 | 34 | 71 | 513          | 91 | 86  | 17  | 48  | 38  | 280     | 793   |

1 "T" refers to Transect number, "Experimental" indicates transects (1-8) in areas where turbines were built between 2003 and 2007, and "Control" indicates transects (9-13) in areas with no turbines in either year.

#### 4.0 DISCUSSION

The results of the grassland bird surveys conducted before and after construction of the windfarm suggest that construction of the facility did not negatively impact numbers of breeding grassland birds. In fact, there was actually an increase in the numbers recorded on the experimental transects (i.e., near the turbines) for six of the nine species analyzed, and overall count totals on the experimental transects were 54% higher after construction (compared to 20% higher on the control transects). O'Connell and Piorkowski (2006) found that of 23 species analyzed at an Oklahoma windfarm, only western meadowlarks were present at lower densities <656 ft (200 m) from turbines than in control areas 0.62 to 6.2 mi (1.0 to 10.0 km) from turbines. The results of other studies (Leddy et al. 1999; Johnson et al. 2000; Erickson et al. 2004) indicated small-scale ( $\leq 164$ -262 ft [50-80 m] from the turbines) decreases in grassland breeding birds. The study design used at JGWEP did not allow for detection of changes at a fine spatial scale, but experimental transects were placed in close proximity to turbines (see Figure 2.3) and numbers of birds detected during surveys increased along those transects after construction.

The 2003 and 2007 surveys were conducted by different personnel, and the increase in birds recorded in 2007 could be attributed to differences in surveyors' abilities to detect birds. However, bird counts on the experimental transects were 54% higher in 2007, and only 20% higher on the 2007 control transects; this suggests an increase in bird numbers on the transects closest to the turbines. The two-sample t-test for total bird numbers did not indicate a significant change, but this is likely due to the high variance between transects in number of birds detected. Multi-year studies, as opposed to the single years pre- and post-construction study conducted at JGWEP, would produce a more robust data set for analysis, and would reduce the probability of a single year potentially having unusually high or low grassland bird numbers (Erickson et al. 2004).

---

## 5.0 LITERATURE CITED

- Erickson, W.P., J. Jeffrey, K. Kronner, and K. Bay. 2004. Stateline Wind project wildlife monitoring final report, July 2001-December 2003. Technical report submitted to FPL Energy, the Oregon Energy Facility Siting Council, and the Stateline Technical Advisory Committee.
- Flath, D.L. 2003. Pre-construction use and mortality of vertebrate wildlife at Judith Gap Wind Resource Area. APEXEnvironmental, Belgrade, Montana.
- Johnson, G.D., W.P. Erickson, M.D. Strickland, M.F. Shepherd, and D.S. Shepherd. 2000. Avian monitoring studies at the Buffalo Ridge, Minnesota Wind Resource Area: Results of a 4-year study. Western Ecosystems Technology, Inc., Cheyenne, Wyoming.
- Leddy, K.L., K.F. Higgins, and D.E. Naugle. 1999. Effects of wind turbines on upland nesting birds in conservation reserve program grasslands. *Wilson Bulletin* 111:100-104.
- O'Connell, T.J., and M.D. Piorkowski. 2006. Sustainable power effects research on wildlife: Final report of 2004-2005 monitoring at the Oklahoma Wind Energy Center. Department of Zoology, Oklahoma State University, Stillwater.

**APPENDIX B:**  
BIRD AND BAT FATALITY TABLES

Table B.1 Bird Fatalities at the Judith Gap Wind Energy Site During the Migration Period (August-October 2006 and February-May 2007).

| Sequential Number | Date                  | Species                | Type of Find <sup>1</sup> | Location/Tower (T) Number    | Distance from Tower (m) | Comments  |
|-------------------|-----------------------|------------------------|---------------------------|------------------------------|-------------------------|---|
| 1                 | 07/05/06 <sup>2</sup> | Western grebe          | IF                        | T18                          | 35 <sup>3</sup>         | Dismembered (head severed; otherwise intact); could be a Clark's grebe, but Western grebe more likely   |
| 2                 | 07/05/06 <sup>2</sup> | Western grebe          | IF                        | T18                          | 35 <sup>3</sup>         | Dismembered (head severed; gash in ventral side); could be Clark's grebe, but Western grebe more likely |
| 3                 | 08/18/06              | Short-eared owl        | CS                        | T2                           | 97                      | Intact carcass  |
| 4                 | 08/21/06              | Unidentified passerine | CS                        | T50                          | 26                      | Intact but old; sparrow or longspur   |
| 5                 | 08/25/06              | Unidentified passerine | CS                        | T32                          | 31                      | Feather spot; 3-4 flight feathers only; possible blackbird  |
| 6                 | 08/29/06              | Horned lark            | CS                        | T34                          | 47                      | Carcass desiccated and scavenged  |
| 7                 | 09/07/06              | Sharp-tailed grouse    | IF                        | T75                          | 7                       | Intact carcass  |
| 8                 | 09/19/06              | American coot          | IF                        | Access road southeast of T99 | 137 <sup>3</sup>        | Intact carcass; trauma to bill and head   |
| 9                 | 09/19/06              | Eared grebe            | CS                        | T32                          | 82                      | Intact carcass; identification tentative; could be a horned grebe                                       |
| 10                | 09/21/06              | Eared grebe            | IF                        | T89                          | 20                      | Intact carcass, but ventral area split open   |
| 11                | 09/21/06              | American coot          | IF                        | Access road near T99         | n/a                     | Intact carcass; trauma to bill and head   |
| 12                | 09/22/06              | Merlin                 | IF                        | By O&M building              | n/a                     | Intact carcass  |
| 13                | 09/23/06              | White-crowned sparrow  | CS                        | T1                           | 113                     | Dismembered (decapitated) and scavenged   |
| 14                | 09/23/06              | Wilson's warbler       | CS                        | T73                          | 86                      | Intact carcass  |
| 15                | 09/23/06              | Wilson's warbler       | CS                        | T73                          | 99                      | Intact carcass  |
| 16                | 10/07/06              | Western meadowlark     | CS                        | T76                          | 17                      | Scavenged carcass   |
| 17                | 10/15/06              | American coot          | CS                        | T99                          | 149                     | Feather spot; identification tentative  |
| 18                | 11/02/06              | American coot          | IF                        | T22                          | 55 <sup>3</sup>         | Intact carcass  |
| 19                | 03/11/06              | Unidentified waterfowl | CS                        | T64                          | 36                      | Feather spot; tegmen visible on flight feather  |

Table B.1 (Continued)

| Sequential Number | Date     | Species     | Type of Find <sup>1</sup> | Location/Tower (T) Number | Distance from Tower (m) | Comments  |
|-------------------|----------|-------------|---------------------------|---------------------------|-------------------------|---|
| 20                | 04/17/07 | Horned lark | CS                        | T83                       | 108                     | Feather spot  |
| 21                | 04/17/07 | Horned lark | CS                        | T83                       | 18                      | Intact carcass  |
| 22                | 05/06/07 | Mallard     | IF                        | T32                       | 11                      | Dismembered (decapitated) and scavenged                     |
| 23                | 05/16/07 | Eared grebe | IF                        | T13                       | 60                      | Intact carcass  |
| 24                | 05/16/07 | Eared grebe | IF                        | T13                       | 55                      | Intact carcass  |
| 25                | 05/26/07 | Eared grebe | CS                        | T31                       | 26                      | Scavenged, but otherwise intact carcass                     |
| 26                | 05/27/07 | Horned lark | CS                        | T2                        | 23                      | Dismembered and heavily scavenged; identification tentative |

<sup>1</sup> Type of find = Carcass was either found during Carcass Search (CS) or was an Incidental Find (IF).

<sup>2</sup> Incidental find prior to initiation of migration study.

<sup>3</sup> Indicates visually estimated distance.

Table B.2 Bat Fatalities at the Judith Gap Wind Energy Site During the Migration Period (August-October 2006 and February-May 2007).

| Sequential Number | Date     | Species               | Type of Find <sup>1</sup> | Location/Tower (T) Number | Distance from Tower (m) | Comments   |
|-------------------|----------|-----------------------|---------------------------|---------------------------|-------------------------|--|
| 1                 | 08/14/06 | Unidentified bat      | CS                        | T6                        | 40                      | Intact carcass   |
| 2                 | 08/14/06 | Hoary bat             | CS                        | T6                        | 11                      | Intact carcass   |
| 3                 | 08/14/06 | Hoary bat             | CS                        | T6                        | 53                      | Intact carcass   |
| 4                 | 08/16/06 | Hoary bat             | CS                        | T1                        | 68                      | Intact carcass   |
| 5                 | 08/16/06 | Hoary bat             | CS                        | T1                        | 62                      | Intact carcass   |
| 6                 | 08/16/06 | Hoary bat             | CS                        | T1                        | 108                     | Intact carcass   |
| 7                 | 08/18/06 | Hoary bat             | CS                        | T2                        | 43                      | Intact carcass   |
| 8                 | 08/18/06 | Hoary bat             | CS                        | T2                        | 13                      | Intact carcass   |
| 9                 | 08/19/06 | Unidentified bat      | CS                        | T99                       | 26                      | Intact carcass   |
| 10                | 08/21/06 | Hoary bat             | CS                        | T50                       | 8                       | Intact carcass   |
| 11                | 08/24/06 | Unidentified bat      | CS                        | T31                       | 36                      | Scavenged carcass; only wings remained                       |
| 12                | 08/25/06 | Hoary bat             | CS                        | T32                       | 51                      | Intact carcass   |
| 13                | 08/25/06 | Unidentified bat      | CS                        | T87                       | 68                      | Scavenged carcass; only wings remained                       |
| 14                | 08/27/06 | Unidentified bat      | CS                        | T63                       | 35                      | Intact carcass   |
| 15                | 08/28/06 | Hoary bat             | CS                        | T73                       | 33                      | Intact carcass but wing scavenged                            |
| 16                | 08/28/06 | Unidentified bat      | CS                        | T73                       | 47                      | Intact carcass but some scavenging or decomposition evident  |
| 17                | 08/28/06 | Hoary bat             | CS                        | T84                       | 31                      | Intact carcass with some scavenging by insects               |
| 18                | 08/28/06 | Hoary bat             | CS                        | T84                       | 14                      | Intact carcass with some decomposition                       |
| 19                | 08/28/06 | Hoary bat             | CS                        | T83                       | 4                       | Intact carcass with some scavenging by insects               |
| 20                | 08/29/06 | Unidentified bat      | CS                        | T40                       | 26                      | Wings only with attached pelage                              |
| 21                | 08/29/06 | Hoary bat             | CS                        | T40                       | 42                      | Wing only with attached pelage                               |
| 22                | 08/29/06 | Silver-haired (?) bat | CS                        | T66                       | 57                      | Intact carcass; tentative identification based on photograph |

Table B.2 (Continued)

| Sequential Number | Date     | Species               | Type of Find <sup>1</sup> | Location/Tower (T) Number | Distance from Tower (m) | Comments   |
|-------------------|----------|-----------------------|---------------------------|---------------------------|-------------------------|--|
| 23                | 08/29/06 | Unidentified bat      | CS                        | T64                       | 23                      | Decomposed carcass   |
| 24                | 08/29/06 | Hoary bat             | CS                        | T64                       | 17                      | Decomposed carcass   |
| 25                | 08/29/06 | Hoary bat             | CS                        | T64                       | 9                       | Carcass intact but decomposed or insect-scavenged            |
| 26                | 09/09/06 | Silver-haired (?) bat | CS                        | T76                       | 72                      | Intact carcass; tentative identification based on photograph |
| 27                | 09/17/06 | Unidentified bat      | CS                        | T40                       | 6                       | Dismembered, scavenged carcass; only wings remained          |
| 28                | 09/23/06 | Unidentified bat      | CS                        | T6                        | 48                      | Scavenged carcass  |
| 29                | 09/23/06 | Unidentified bat      | CS                        | T73                       | 54                      | Scavenged carcass; only parts of wings remained              |
| 30                | 10/07/06 | Unidentified bat      | IF                        | T42                       | 33                      | Only scavenged wing remained                                 |
| 31                | 10/09/06 | Unidentified bat      | CS                        | T66                       | 16                      | Dismembered, insect scavenged carcass                        |
| 32                | 10/09/06 | Unidentified bat      | CS                        | T66                       | 17                      | Only scavenged wings remained                                |
| 33                | 10/13/06 | Silver-haired (?) bat | CS                        | T34                       | 10                      | Intact carcass; tentative identification based on photograph |
| 34                | 10/14/06 | Unidentified bat      | CS                        | T83                       | 108                     | Intact but decomposed carcass                                |
| 35                | 10/15/06 | Hoary bat             | CS                        | T73                       | 100                     | Scavenged, decomposed carcass                                |
| 36                | 05/19/07 | Silver-haired bat     | CS                        | T63                       | 50                      | Intact carcass   |

<sup>1</sup> Type of find = Carcass Search (CS) or Incidental Find (IF).

**APPENDIX C:**  
FORMS

**JUDITH GAP WIND ENERGY PROJECT  
FATALITY/INJURED WILDLIFE DATA FORM**

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_ OBSERVER: \_\_\_\_\_

FOUND DURING (check one): SCHEDULED CARCASS SEARCH \_\_\_\_\_ INCIDENTAL FIND \_\_\_\_\_

COLLECTED? yes no ID<sup>1</sup>: \_\_\_\_\_ PHOTO NOS: \_\_\_\_\_



WHEN TAKING THE PHOTOGRAPHS, INCLUDE AN OBJECT FOR SCALE, PHOTOGRAPH THE BACK AND THE UNDERSIDE, AND INCLUDE CLOSE-UPS OF HEAD, FEET, AND PLUMAGE.

PLOT NO.: \_\_\_\_\_

LOCATION, IF NOT ON PLOT: \_\_\_\_\_

HABITAT: \_\_\_\_\_

SPECIES: \_\_\_\_\_ SEX (circle): M F U AGE (circle): A J U

CONDITION (circle one): intact scavenged dismembered feather spot injured

**DISTANCE & BEARING FROM THE NEAREST TOWER/POLE:**

DESCRIPTION DISTANCE (ft) BEARING (degrees)

Part 1 \_\_\_\_\_

Part 2 \_\_\_\_\_

Part 3 \_\_\_\_\_

Other \_\_\_\_\_

COMMENTS:

ESTIMATED TIME SINCE DEATH/INJURY: \_\_\_\_\_

**WEATHER:**

HISTORY: (if carcass is estimated to be less than one week old, circle any of the following weather conditions that occurred at or before the estimated time of death/incident)

clear calm fog cloudy rain snow storm gusty wind violent storm blizzard

NOTES:

**GENERAL COMMENTS:** (e.g., behavior observed if bird is injured; details of carcass – body parts missing, injuries, number of feathers in feather spot; indications of cause of death, field marks for identification, USFWS band no., etc.)

**USFWS CONTACT:**

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_ RECOVERY APPROVAL: yes no

CONTACT PERSON(S):

COMMENTS:

DISPOSITION OF FIND: \_\_\_\_\_

TRANSPORTED TO FREEZER: \_\_\_\_\_ DATE: \_\_\_\_\_ TIME: \_\_\_\_\_

RELEASE TO USFWS - PERSON: \_\_\_\_\_ DATE: \_\_\_\_\_ TIME: \_\_\_\_\_

COMMENTS:


<sup>1</sup> ID = mm-dd-yy-hrmin (e.g., 08-24-06-1307).





{Nest Number} Photo Page

{Nest Number} Photo Page



Photograph 1 {Nest Number}, Looking \_\_\_\_\_.





