

Circular Air-Filled Space in the Aortic Knob:

A Rare Radiologic Finding

Magdy M. El-Sayed Ahmed,
MD, MSc
Muhammad Aftab, MD
Kim I. de la Cruz, MD
Reynolds M. Delgado III, MD
Hari R. Mallidi, MD

Section Editor:

Raymond F. Stainback, MD,
Department of Adult
Cardiology, Texas Heart
Institute, 6624 Fannin St.,
Suite 2480, Houston, TX
77030

From: Departments of Cardiovascular Surgery (Drs. Aftab, de la Cruz, El-Sayed Ahmed, and Mallidi) and Cardiology (Dr. Delgado), Texas Heart Institute, Houston, Texas 77030; Division of Cardiothoracic Surgery, Michael E. DeBakey Department of Surgery (Drs. Aftab, de la Cruz, and Mallidi), Baylor College of Medicine, Houston, Texas 77030; and Department of Surgery (Dr. El-Sayed Ahmed), Faculty of Medicine, Zagazig University, Zagazig 44519, Egypt

Address for reprints:

Magdy M. El-Sayed Ahmed,
MD, MSc, Department of
Cardiovascular Surgery,
WW 3 Suite 400, Children's
National Medical Center,
111 Michigan Ave. NW,
Washington, DC 20010

E-mail: elgoharymagdy@
yahoo.com

© 2014 by the Texas Heart®
Institute, Houston

A 70-year-old man with a history of orthotopic heart transplantation developed severe coronary allograft vasculopathy and presented with cardiogenic shock. He was resuscitated, and both inotropic and vasopressor support were initiated. Subsequently, an intra-aortic balloon pump (IABP) was inserted for further hemodynamic support. After 4 weeks, the IABP was moved from the common femoral artery to the left axillary artery, to motivate the patient to exercise and enable him to sit up in bed. The procedure was performed in the hybrid suite under fluoroscopic guidance. The left axillary artery was surgically exposed, and a 6-mm Vascutek® Gelweave™ graft (Vascutek Ltd., a Terumo company; Renfrewshire, Scotland) was anastomosed as a conduit. Routine postoperative subclavian IABP management included twice-weekly chest radiographs to monitor for possible device migration, and daily examination of the patient's left hand and left radial artery pulse. A portable chest radiograph in the anteroposterior view revealed an unusual, circular, air-filled space in the aortic knob (transverse aorta) (Fig. 1). This finding resulted from the coincidental "end-on" image acquisition of the inflated and malpositioned IABP that had migrated proximally to the transverse aorta.

Comment

The IABP is frequently used for mechanical support in cardiac procedures.¹ Transfemoral artery insertion, the traditional approach, might be contraindicated in patients with severe atherosclerosis of the iliofemoral arteries or the aorta, and in patients with aortobifemoral bypass grafts. Alternative approaches are insertion through the transaxillary artery,² the transbrachial artery,^{3,4} or through grafts to the ascending aorta.⁵ Mayer⁶ was the first to describe the trans-subclavian artery insertion of an IABP; McBride and colleagues⁷ subsequently described a simplified technique.

This radiologic image conveys a teaching and educational message of a very rare finding that was captured incidentally. The finding is unusual because the balloon, while turning from the left subclavian artery into the descending thoracic aorta, displayed an anteroposterior curve, creating a circular air-filled space in the aortic knob—a cross-section of the balloon, captured during inflation. This chest radio-




Fig. 1 Chest radiograph (anteroposterior view) shows a circular air-filled space in the aortic knob (arrow). Arrowheads point to the intra-aortic balloon pump radiopaque markers.

graph shows an important sign of inappropriate IABP positioning in the transverse aorta that would be captured only upon the coincidence of image acquisition during balloon inflation. This image would not have been acquired if the IABP had been appropriately placed in the descending thoracic aorta distal to the origin of the left subclavian artery, so that the IABP contour remained relatively straight.

References

1. Christenson JT, Buswell L, Velebit V, Maurice J, Simonet F, Schmuziger M. The intraaortic balloon pump for postcardiotomy heart failure. Experience with 169 intraaortic balloon pumps. *Thorac Cardiovasc Surg* 1995;43(3):129-33.
2. H'Doubler PB Jr, H'Doubler WZ, Bien RC, Jansen DA. A novel technique for intraaortic balloon pump placement via the left axillary artery in patients awaiting cardiac transplantation. *Cardiovasc Surg* 2000;8(6):463-5.
3. Onorati F, Impiombato B, Ferraro A, Comi MC, Spaccarotella C, Indolfi C, Renzulli A. Transbrachial intraaortic balloon pumping in severe peripheral atherosclerosis. *Ann Thorac Surg* 2007;84(1):264-6.
4. Bundhoo S, O'Keefe PA, Luckraz H, Ossei-Gerning N. Extended duration of brachially inserted intra-aortic balloon pump for myocardial protection in two patients undergoing urgent coronary artery bypass grafting. *Interact Cardiovasc Thorac Surg* 2008;7(1):42-4.
5. Burack JH, Uceda P, Cunningham JN Jr. Transthoracic intraaortic balloon pump: a simplified technique. *Ann Thorac Surg* 1996;62(1):299-301.
6. Mayer JH. Subclavian artery approach for insertion of intra-aortic balloon. *J Thorac Cardiovasc Surg* 1978;76(1):61-3.
7. McBride LR, Miller LW, Naunheim KS, Pennington DG. Axillary artery insertion of an intraaortic balloon pump. *Ann Thorac Surg* 1989;48(6):874-5.